



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.


About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





27-740



5.2-1827
SOLUTIONS

OF

THE MORE DIFFICULT EQUATIONS

CONTAINED

IN THE FOURTH EDITION

OF

DR. BLAND'S ALGEBRAICAL PROBLEMS.

BY

FRANCIS EDWARD THOMPSON, B.A.

OF TRINITY COLLEGE, CAMBRIDGE.

PA. Jube SOLVI, obsecro.
Sr. Age, sat. PA. At matura. Sr. Eo intrō.
PA. O faustum, et felicem, hanc diem !—T.E.

LONDON:

PRINTED FOR J. MAWMAN, LUDGATE-STREET.

MDCCCXXVII.

740



LONDON
Printed by J. CLOWES,
Stationer-street.

ADVERTISEMENT.

IN publishing the following Solutions, the Author has no intention of producing indolence in the learner. He has found that the beginner, who has no tutor to consult, and is compelled to depend upon his own resources, is often discouraged at the very commencement of his attempts. In fact, the solution of the majority of the following Equations consists in some trick—when that is discovered, all is finished. *Ce n'est que le premier pas qui coute.* But to take this first step is the very difficulty—and it is the object of the following pages to remove it.

The Author has given Solutions only for the last few of the Simple Equations involving one unknown Quantity; they appear to be the only ones requiring explanation.

The Simple Equations involving two unknown Quantities are omitted altogether—they may all be solved by application of the ordinary rules.

The Pure Quadratics are all solved. Solutions of the Affected Quadratics involving one unknown Quantity

will be found for all that are really difficult ; and similarly for those involving two unknown quantities.

The Reader will thus perceive that it has been the Author's aim to produce a compendious and useful manual. He might have swelled it out by giving Solutions to all the Equations—but this was far from his design.

Should this tract be received with approbation, it is the Author's intention to prepare Solutions of the more difficult Problems contained in the same work.

SOLUTIONS

OF

SIMPLE AND QUADRATIC EQUATIONS.

Solutions of Simple Equations involving one unknown Quantity.

Bland, p. 299.

46. Given $\sqrt[3]{10x + 35} - 1 = 4$, to find the value of x .

Transposing $\sqrt[3]{10x + 35} = 5$,

and cubing each side, $10x + 35 = 125$;

$\therefore 10x = 90$,

and $x = 9$.

47. Given $\sqrt[5]{9x - 4} + 6 = 8$, to find the value of x .

By transposition $\sqrt[5]{9x - 4} = 2$,

and involving each side to the fifth power,

$9x - 4 = 32$;

$\therefore 9x = 36$,

and $x = 4$.

48. Given $\sqrt{x + 16} = 2 + \sqrt{x}$, to find the value of x .

Squaring both sides, $x + 16 = 4 + 4\sqrt{x} + x$;

$\therefore 4\sqrt{x} = 12$,

$\sqrt{x} = 3$,

and $x = 9$.

49. Given $\sqrt{x-32} = 16 - \sqrt{x}$, to find the value of x .

Squaring both sides, $x - 32 = 256 - 32\sqrt{x} + x$;

$$\therefore 32\sqrt{x} = 288,$$

$$\sqrt{x} = 9,$$

$$\text{and } x = 81.$$

50. Given $\sqrt{4x+21} = 2\sqrt{x} + 1$, to find the value of x .

Squaring each side, $4x + 21 = 4x + 4\sqrt{x} + 1$;

$$\therefore 4\sqrt{x} = 20,$$

$$\sqrt{x} = 5,$$

$$\text{and } x = 25.$$

51. Given $a\sqrt[3]{bx-c} = d\sqrt[3]{ex+fx-g}$, to find the value of x .

Cubing each side, $a^3(bx-c) = d^3(ex+fx-g)$,

$$\text{or } a^3bx - a^3c = d^3ex + d^3fx - d^3g;$$

by transposition, $a^3bx - d^3ex - d^3fx = a^3c - d^3g$,

$$\text{or } x(a^3b - d^3(e+f)) = a^3c - d^3g;$$

$$\therefore x = \frac{a^3c - d^3g}{a^3b - d^3(e+f)}.$$

52. Given $\sqrt[3]{a^3+c} = \sqrt[4]{\frac{a^3+c}{d(x+b)}}$, to find the value of x .

Involving each side to the 12th power,

$$(a^3+c)^4 = \left(\frac{a^3+c}{d(x+b)}\right)^3,$$

$$\text{or } (d(x+b))^3 = \frac{1}{a^3+c};$$

Extracting the cube-root on both sides,

$$\begin{aligned} d \cdot \overline{x+b} &= \frac{1}{\sqrt[3]{a^3+c}}; \\ \therefore x+b &= \frac{1}{d \cdot \sqrt[3]{a^3+c}}, \\ \text{and } x &= \frac{1}{d \cdot \sqrt[3]{a^3+c}} - b. \end{aligned}$$

53. Given $\sqrt[m]{a+x} = \sqrt[m]{x^2+5ax+b^2}$, to find the value of x .

Raising both sides to the $2m^{\text{th}}$ power,

$$\begin{aligned} a^2 + 2ax + x^2 &= x^2 + 5ax + b^2, \\ \text{or } 3ax &= a^2 - b^2; \\ \therefore x &= \frac{a^2 - b^2}{3a}. \end{aligned}$$

54. Given $a+b \cdot \sqrt[m]{x+d} = c$.

$$\text{Now } \sqrt[m]{x+d} = \frac{c-a}{b};$$

$$\therefore x+d = \left(\frac{c-a}{b} \right)^m,$$

$$\text{and } x = \left(\frac{c-a}{b} \right)^m - d.$$

55. Given $\frac{\sqrt{9x}-4}{\sqrt{x}+2} = \frac{15+\sqrt{9x}}{\sqrt{x}+40}$, to find the value of x .

Clearing of fractions, we have,

$$3x + 116\sqrt{x} - 160 = 3x + 21\sqrt{x} + 30;$$

$$\text{then } 95\sqrt{x} = 190,$$

$$\sqrt{x} = 2,$$

$$\text{and } x = 4.$$

56. Given $\frac{\sqrt{x} + \sqrt{b}}{\sqrt{x} - \sqrt{b}} = \frac{a}{b}$, to find the value of x .

This may be solved by two methods—1st, by the common rules of solution; and 2ndly, by the assistance of proportion.

$$\text{1st, then, } b\sqrt{x} + b\sqrt{b} = a\sqrt{x} - a\sqrt{b},$$

$$\text{or } a\sqrt{x} - b\sqrt{x} = a\sqrt{b} + b\sqrt{b};$$

$$\therefore \sqrt{x} = \sqrt{b} \left(\frac{a+b}{a-b} \right),$$

$$\text{and } x = b \cdot \left(\frac{a+b}{a-b} \right)^2.$$

2ndly, more neatly thus:

$$\sqrt{x} + \sqrt{b} : \sqrt{x} - \sqrt{b} :: a : b,$$

$$\text{or } \sqrt{x} : \sqrt{b} :: a+b : a-b, \text{ (Wood's Alg., 182.)}$$

$$\therefore \sqrt{x} = \sqrt{b} \cdot \left(\frac{a+b}{a-b} \right),$$

$$\text{and } x = b \cdot \left(\frac{a+b}{a-b} \right)^2.$$

57. Given $\frac{\sqrt{6x}-2}{\sqrt{6x}+2} = \frac{4\sqrt{6x}-9}{4\sqrt{6x}+6}$, to find the value of x .

$$\text{Now } \sqrt{6x} + 2 : \sqrt{6x} - 2 :: 4\sqrt{6x} + 6 : 4\sqrt{6x} - 9,$$

$$\text{or } \sqrt{6x} : 2 :: 8\sqrt{6x} - 3 : 15 \quad (\text{Alg. 182});$$

$$\therefore 15\sqrt{6x} = 16\sqrt{6x} - 6,$$

$$\sqrt{6x} = 6,$$

$$6x = 36,$$

$$\text{and } x = 6.$$

58. Given $\frac{5x-9}{\sqrt{5x}+3} - 1 = \frac{\sqrt{5x}-3}{2}$, to find the value of x .

$$\text{Then } (\sqrt{5x} - 3) - 1 = \frac{\sqrt{5x} - 3}{2},$$

$$\text{or } \frac{\sqrt{5x} - 3}{2} = 1;$$

$$\therefore \sqrt{5x} - 3 = 2,$$

$$\sqrt{5x} = 5,$$

$$5x = 25,$$

$$\text{and } x = 5.$$

59. Given $\sqrt{1+x}\sqrt{x^2+12} = 1+x$, to find the value of x ,
squaring both sides of the equation.

$$1 + x\sqrt{x^2+12} = 1 + 2x + x^2,$$

$$\text{and } \sqrt{x^2+12} = x + 2 \quad \text{or } x=0$$

then squaring each side, $x^2 + 12 = x^2 + 4x + 4$.

$$\therefore 4x = 8$$

$$\text{and } x = 2$$

60. Given $\frac{ax}{b} \cdot \sqrt{c^2x^2 + d^2} + \frac{acx^2}{b} = ex$, to find the
value of x .

Dividing both sides by $\frac{x}{b}$,

$$a\sqrt{c^2x^2 + d^2} + acx = eb \quad \text{or } x=0$$

by transposition $a\sqrt{c^2x^2 + d^2} = eb - acx$,

squaring both sides, $a^2c^2x^2 + a^2d^2 = e^2b^2 - 2abcex + a^2c^2x^2$.

$$\therefore 2abcex = e^2b^2 - a^2d^2$$

$$\text{and } x = \frac{b^2e^2 - a^2d^2}{2abce}$$

61. Given $\sqrt{x} + \sqrt{x-9} = \frac{36}{\sqrt{x-9}}$, to find the value
of x .

Clearing of fractions,

$$\sqrt{x^2 - 9x} + x - 9 = 36$$

$$\text{and } \sqrt{x^2 - 9x} = 45 - x$$

squaring both sides, $x^2 - 9x = 2025 - 90x + x^2$.

$$\therefore 81x = 2025$$

$$\text{and } x = 25.$$

Solutions of Pure Quadratics and others, &c.

1. Given $3x^2 - 4 = 28 + x^2$, to find the values of x .

$$\therefore 2x^2 = 32$$

$$x^2 = 16$$

$$\text{and } x = \pm 4$$

2. Given $x + y : y :: 3 : 1$ } to find the values of x
and $xy = 18$ } and y .

From the first equation, $x : y :: 2 : 1$ (*Alg.* 180.)

$$\therefore x = 2y$$

$$\text{then by substitution } 2y^2 = 18$$

$$\therefore y^2 = 9$$

$$\text{and } y = \pm 3$$

3. Given $x - y : y :: 4 : 5$ } to find the values of x
and $x^2 + 4y^2 = 181$ } and y .

From the first equation, $x : y :: 9 : 5$. (*Alg.* 179.)

$$\text{and } x = \frac{9y}{5}$$

substituting this value in the second equation,

$$\frac{81y^2}{25} + 4y^2 = 181$$

$$\therefore 81y^2 + 100y^2 = 25 \times 181$$

$$y^2 = 25$$

$$\text{and } y = \pm 5$$

$$\text{and } x = \frac{9y}{5} = \pm 9.$$

$$4. \text{ Given } x + y : x - y :: a : b \left. \vphantom{\begin{matrix} x + y \\ x - y \end{matrix}} \right\} \text{ to find the values of } x \text{ and } y.$$

$$\text{and } xy = c^2$$

From the first equation, $x : y :: a + b : a - b$. (Alg. 182.)

$$\text{and } x = y \cdot \frac{a+b}{a-b}$$

Then substituting this value in the second equation,

$$y^2 \cdot \frac{a+b}{a-b} = c^2$$

$$\text{and } y^2 = c^2 \cdot \frac{a-b}{a+b},$$

$$y = \pm c \sqrt{\frac{a-b}{a+b}},$$

$$\text{and } x = y \cdot \frac{a+b}{a-b} = \pm c \cdot \sqrt{\frac{a+b}{a-b}}.$$

$$5. \text{ Given } x^2 + y^2 : x^2 - y^2 :: 17 : 8 \left. \vphantom{\begin{matrix} x^2 + y^2 \\ x^2 - y^2 \end{matrix}} \right\} \text{ to find the values of } x \text{ and } y.$$

$$\text{and } xy^2 = 45$$

From the first equation, $x^2 : y^2 :: 25 : 9$. (Alg. 182.)

$$\text{and } x : y :: 5 : 3. \text{ (Alg. 188.)}$$

$$\therefore x = \frac{5y}{3},$$

substituting this value in the second equation,

$$\frac{5y^3}{3} = 45;$$

$$\therefore y^3 = 27,$$

$$\text{and } y = 3;$$

$$\therefore x = \frac{5y}{3} = 5.$$

Solutions of Pure Quadratics.

6. Given $x^2 - xy = 54$ } to find the values of x and
 and $xy - y^2 = 18$ } y .

$$\text{by subtraction, } x^2 - 2xy + y^2 = 36,$$

$$\text{extracting the square root } x - y = \pm 6.$$

$$\text{Then from the first equation, } \therefore x \cdot \overline{x-y} = 54,$$

$$\text{and } x = \frac{54}{\pm 6} = \pm 9;$$

$$\text{and from the second equation, } y \cdot \overline{x-y} = 18;$$

$$\therefore y = \frac{18}{\pm 6} = \pm 3.$$

7. Given $x + y : x^2 - y^2 :: 1 : 4$ } to find the values of
 and $xy = 21$ } x and y .

Dividing the two first terms of the proportion by x and y .

$$1 : x - y :: 1 : 4;$$

$$\therefore x - y = 4,$$

$$\text{squaring both sides, } x^2 - 2xy + y^2 = 16,$$

$$\text{but } 4xy = 84$$

$$\text{by addition, } x^2 + 2xy + y^2 = 100;$$

$$\therefore x + y = \pm 10,$$

$$\text{but } x - y = 4;$$

$$\therefore x = 7 \text{ or } -3;$$

$$\text{and } y = 3 \text{ or } -7.$$

8. Given $ax^2 + bxy = c^2$ } to find the values of x
 and $x-y : x :: m : n$ } and y .

$$\text{From the second equation, } y : x :: n - m : n;$$

$$\therefore y = x \cdot \frac{n-m}{n},$$

Substituting this value in the first equation,

$$ax^2 + bx^2 \cdot \frac{n-m}{n} = c^2;$$

$$\therefore x^2 \left(\frac{na + nb - mb}{n} \right) = c^2;$$

$$\text{and } x^2 = \frac{nc^2}{na + nb - mb};$$

$$\therefore x = \pm c \cdot \sqrt{\frac{nc}{na + nb - mb}}.$$

$$\text{and } y = \frac{x \cdot \overline{n-m}}{n} = \pm \frac{n-m}{n} \cdot c \cdot \sqrt{\frac{nc}{na + nb - mb}}.$$

9. Given $x^2 + y^2 : x^2 - y^2 :: 559 : 127$ } to find the values
and $x^2 y = 294$ } of x and y .

From the first equation, $2x^2 : 2y^2 :: 686 : 432$. (*Alg.* 182.)

$$\text{or } x^2 : y^2 :: 343 : 216;$$

$$\text{or } x : y :: 7 : 6. \text{ (*Alg.* 188.)};$$

$$\text{and } y = \frac{6x}{7}.$$

$$\text{Now } \frac{6x^2}{7} = 294;$$

$$\text{or } \frac{x^2}{7} = 49;$$

$$\text{then } x^2 = 343;$$

$$\therefore x = 7,$$

$$\text{and } y = \frac{6x}{7} = 6.$$

10. Given $x^2 - xy : xy - y^2 :: 3 : 7$ } to find the values
and $xy^2 = 147$ } of x and y .

Dividing the proportion by $x - y$ we have,

$$x : y :: 3 : 7;$$

$$\therefore x = \frac{3y}{7}.$$

Substituting this value in the second equation,

$$\frac{3y^2}{7} = 147;$$

$$\therefore \frac{y^2}{7} = 49;$$

$$y^2 = 343,$$

$$\text{and } y = 7;$$

$$\therefore x = \frac{3y}{7} = 3.$$

11. Given $\sqrt{x} + \sqrt{y} : \sqrt{x} - \sqrt{y} :: 4 : 1$ } to find
 and $x - y = 16$
 the values of x and y .

From the first equation, $\sqrt{x} : \sqrt{y} :: 5 : 3$. (*Alg.* 182.)

$$\text{and } \sqrt{x} = \frac{5\sqrt{y}}{3};$$

$$\therefore x = \frac{25y}{9}.$$

Substituting this value in the second equation,

$$\frac{25y}{9} - y = 16,$$

$$25y - 9y = 144;$$

$$\therefore y = \frac{144}{16} = 9,$$

$$\text{and } x = \frac{25y}{9} = 25.$$

12. Given $\sqrt[4]{x} - \sqrt[4]{y} = 3$ } to find the values of x and y .
 and $\sqrt[4]{x} + \sqrt[4]{y} = 7$ }

$$\therefore \sqrt[4]{x} = 5;$$

$$\text{and } x = 625.$$

$$\begin{aligned}\text{Again, } \sqrt[4]{y} &= 2, \\ y &= 16.\end{aligned}$$

$$\begin{aligned}13. \text{ Given } x - y : \sqrt{x} - \sqrt{y} :: 8 : 1 \\ \text{and } \sqrt{xy} = 15\end{aligned} \left. \vphantom{\begin{aligned} x - y : \sqrt{x} - \sqrt{y} :: 8 : 1 \\ \text{and } \sqrt{xy} = 15 \end{aligned}} \right\} \text{to find the values of } x \text{ and } y.$$

Dividing the first equation by $\sqrt{x} - \sqrt{y}$;

$$\therefore \sqrt{x} + \sqrt{y} : 1 :: 8 : 1,$$

$$\text{and } \sqrt{x} + \sqrt{y} = 8;$$

$$\text{squaring each side } x + 2\sqrt{xy} + y = 64;$$

$$\text{but } 4\sqrt{xy} = 60;$$

$$\therefore x - 2\sqrt{xy} + y = 4,$$

$$\text{and } \sqrt{x} - \sqrt{y} = \pm 2;$$

$$\text{but } \sqrt{x} + \sqrt{y} = 8;$$

$$\therefore \sqrt{x} = 5 \text{ or } 3,$$

$$x = 25 \text{ or } 9;$$

$$\text{and } \sqrt{y} = 3 \text{ or } 5;$$

$$\text{and } y = 9 \text{ or } 25.$$

$$\begin{aligned}14. \text{ Given } x^3 - y^3 : x^2y - xy^2 :: 7 : 2 \\ \text{and } x + y = 6\end{aligned} \left. \vphantom{\begin{aligned} x^3 - y^3 : x^2y - xy^2 :: 7 : 2 \\ \text{and } x + y = 6 \end{aligned}} \right\} \text{to find the values of } x \text{ and } y.$$

From the first equation, $x^3 - y^3 : 3x^2y - 3xy^2 :: 7 : 6$;

$$\text{and } x^3 - 3x^2y + 3xy^2 - y^3 : x^2y - xy^2 :: 1 : 2. \text{ (Alg. 180.)}$$

$$\text{or } (x - y)^3 : xy.(x - y) :: 1 : 2;$$

$$\therefore (x - y)^2 : xy :: 1 : 2;$$

$$\text{and } x^2 - 2xy + y^2 : 4xy :: 1 : 8;$$

$$\therefore x^2 - 2xy + y^2 : x^2 + 2xy + y^2 :: 1 : 9,$$

$$x - y : x + y :: 1 : \pm 3;$$

$$\text{and } x - y : 6 :: 1 : \pm 3;$$

Substituting this value in the second equation,

$$\frac{3y^2}{7} = 147;$$

$$\therefore \frac{y^2}{7} = 49;$$

$$y^2 = 343,$$

$$\text{and } y = 7;$$

$$\therefore x = \frac{3y}{7} = 3.$$

11. Given $\sqrt{x} + \sqrt{y} : \sqrt{x} - \sqrt{y} :: 4 : 1$ } to find
and $x - y = 16$ }
the values of x and y .

From the first equation, $\sqrt{x} : \sqrt{y} :: 5 : 3$. (*Alg.* 182.)

$$\text{and } \sqrt{x} = \frac{5\sqrt{y}}{3};$$

$$\therefore x = \frac{25y}{9}.$$

Substituting this value in the second equation,

$$\frac{25y}{9} - y = 16,$$

$$25y - 9y = 144;$$

$$\therefore y = \frac{144}{16} = 9,$$

$$\text{and } x = \frac{25y}{9} = 25.$$

12. Given $\sqrt[4]{x} - \sqrt[4]{y} = 3$ } to find the values of x and y .
and $\sqrt[4]{x} + \sqrt[4]{y} = 7$ }

$$\therefore \sqrt[4]{x} = 5;$$

$$\text{and } x = 625.$$

$$\begin{aligned}\text{Again, } \sqrt[4]{y} &= 2, \\ y &= 16.\end{aligned}$$

$$\begin{aligned}13. \quad \text{Given } x - y : \sqrt{x} - \sqrt{y} :: 8 : 1 \\ \text{and } \sqrt{xy} = 15\end{aligned} \left. \vphantom{\begin{aligned} \text{Given } x - y : \sqrt{x} - \sqrt{y} :: 8 : 1 \\ \text{and } \sqrt{xy} = 15 \end{aligned}} \right\} \text{to find the values of } x \text{ and } y.$$

Dividing the first equation by $\sqrt{x} - \sqrt{y}$;

$$\therefore \sqrt{x} + \sqrt{y} : 1 :: 8 : 1,$$

$$\text{and } \sqrt{x} + \sqrt{y} = 8;$$

$$\text{squaring each side } x + 2\sqrt{xy} + y = 64;$$

$$\text{but } 4\sqrt{xy} = 60;$$

$$\therefore x - 2\sqrt{xy} + y = 4,$$

$$\text{and } \sqrt{x} - \sqrt{y} = \pm 2;$$

$$\text{but } \sqrt{x} + \sqrt{y} = 8;$$

$$\therefore \sqrt{x} = 5 \text{ or } 3,$$

$$x = 25 \text{ or } 9;$$

$$\text{and } \sqrt{y} = 3 \text{ or } 5;$$

$$\text{and } y = 9 \text{ or } 25.$$

$$\begin{aligned}14. \quad \text{Given } x^3 - y^3 : x^2y - xy^2 :: 7 : 2 \\ \text{and } x + y = 6\end{aligned} \left. \vphantom{\begin{aligned} \text{Given } x^3 - y^3 : x^2y - xy^2 :: 7 : 2 \\ \text{and } x + y = 6 \end{aligned}} \right\} \text{to find the values of } x \text{ and } y.$$

From the first equation, $x^3 - y^3 : 3x^2y - 3xy^2 :: 7 : 6$;

and $x^3 - 3x^2y + 3xy^2 - y^3 : x^2y - xy^2 :: 1 : 2$. (*Alg.* 180.)

$$\text{or } (x - y)^3 : xy.(x - y) :: 1 : 2;$$

$$\therefore (x - y)^2 : xy :: 1 : 2;$$

$$\text{and } x^2 - 2xy + y^2 : 4xy :: 1 : 8;$$

$$\therefore x^2 - 2xy + y^2 : x^2 + 2xy + y^2 :: 1 : 9,$$

$$x - y : x + y :: 1 : \pm 3;$$

$$\text{and } x - y : 6 :: 1 : \pm 3;$$

$$\begin{aligned}\therefore x - y &= \pm 2, \\ \text{and } x + y &= 6.\end{aligned}$$

$$\begin{aligned}\therefore x &= 4 \text{ or } 2, \\ \text{and } y &= 2 \text{ or } 4.\end{aligned}$$

$$15. \quad \left. \begin{aligned} \text{Given } \frac{1}{x} + \frac{1}{y} &= \frac{1}{2} \\ \text{and } \frac{2}{xy} &= \frac{1}{9} \end{aligned} \right\} \text{to find the values of } x \text{ and } y.$$

From the first equation, $x + y = \frac{xy}{2}$;

$$\text{and } x + y = \frac{18}{2} = 9,$$

$$\begin{aligned}\therefore x^2 + 2xy + y^2 &= 81, \\ \text{but } 4xy &= 72.\end{aligned}$$

by subtraction, $x^2 - 2xy + y^2 = 9$;

$$\therefore x - y = \pm 3,$$

$$\text{and } x + y = 9;$$

$$\therefore x = 6 \text{ or } 3;$$

$$\text{and } y = 3 \text{ or } 6.$$

$$16. \quad \left. \begin{aligned} \text{Given } x^4 - y^4 &= 369 \\ \text{and } x^2 - y^2 &= 9 \end{aligned} \right\} \text{to find the values of } x \text{ and } y.$$

Dividing the first equation by the second,

$$x^2 + y^2 = 41,$$

$$\text{but } x^2 - y^2 = 9.$$

$$\therefore x^2 = 25,$$

$$\text{and } x = \pm 5,$$

$$y^2 = 16;$$

$$\text{and } y = \pm 4.$$

$$17. \quad \left. \begin{array}{l} \text{Given } x^3 - y^3 = 56, \\ \text{and } x - y = \frac{16}{xy} \end{array} \right\} \text{to find the values of } x \text{ and } y.$$

$$\begin{array}{l} x^3 - y^3 = 56, \\ \text{and } 3x^2y - 3xy^2 = 48 \text{ (from the 2nd equation);} \end{array}$$

\therefore by subtraction, $x^3 - 3x^2y + 3xy^2 - y^3 = 8$,
and extracting the cube-root on both sides,

$$\begin{array}{l} x - y = 2; \\ \therefore x^3 - 2xy + y^3 = 4, \\ \text{but } 4xy = 32; \end{array}$$

$$\begin{array}{l} \therefore x^3 + 2xy + y^3 = 36, \\ \text{and } x + y = \pm 6; \\ \text{but } x - y = 2. \end{array}$$

$$\begin{array}{l} \therefore x = 4 \text{ or } -2, \\ \text{and } y = 2 \text{ or } -4. \end{array}$$

18. Given $\frac{1}{1-\sqrt{1-x^2}} - \frac{1}{1+\sqrt{1-x^2}} = \frac{\sqrt{3}}{x^2}$, to find
the values of x .

Clearing the equation of fractions,

$$1 + \sqrt{1-x^2} - 1 + \sqrt{1-x^2} = (1 - \sqrt{1-x^2})(1 + \sqrt{1-x^2}) \cdot \frac{\sqrt{3}}{x^2},$$

$$\text{or } 2\sqrt{1-x^2} = x^2 \cdot \frac{\sqrt{3}}{x^2},$$

$$\text{or } 2\sqrt{1-x^2} = \sqrt{3},$$

$$\text{or } \sqrt{1-x^2} = \frac{\sqrt{3}}{2};$$

$$\therefore 1 - x^2 = \frac{3}{4},$$

$$x^2 = \frac{1}{4},$$

$$\text{and } x = \pm \frac{1}{2}.$$

$$19. \left. \begin{array}{l} x^2y + y^3 = 116 \\ \text{and } xy^{\frac{1}{2}} + y = 14 \end{array} \right\} \text{to find the values of } x \text{ and } y.$$

Squaring the second equation, we have

$$\begin{array}{r} x^2y + 2xy^{\frac{3}{2}} + y^3 = 196 \\ \text{but } x^2y \qquad \qquad + y^3 = 116; \\ \hline \end{array}$$

$$\therefore 2xy^{\frac{3}{2}} = 80,$$

$$\text{and } x^2y - 2xy^{\frac{3}{2}} + y^3 = 36;$$

$$\therefore xy^{\frac{1}{2}} - y = \pm 6,$$

$$\text{but } xy^{\frac{1}{2}} + y = 14;$$

$$\therefore y = 4 \text{ or } 10,$$

$$\text{and } xy^{\frac{1}{2}} = 10 \text{ or } 4;$$

$$\therefore x = \frac{10}{y^{\frac{1}{2}}} \text{ or } \frac{4}{y^{\frac{1}{2}}},$$

$$= \frac{10}{2}, \text{ or } \frac{4}{\sqrt{10}},$$

$$= 5, \text{ or } \sqrt{\frac{16}{10}},$$

$$= 5, \text{ or } 2\sqrt{\frac{2}{5}}.$$

$$20. \left. \begin{array}{l} \sqrt[3]{x} + \sqrt[3]{y} = 6, \\ \text{and } x + y = 72, \end{array} \right\} \text{to find the values of } x \text{ and } y.$$

Cubing the first equation, we have

$$\begin{array}{r} x + 3x^{\frac{2}{3}}y^{\frac{1}{3}} + 3x^{\frac{1}{3}}y^{\frac{2}{3}} + y = 216, \\ \text{but } x \qquad \qquad \qquad + y = 72; \\ \hline \end{array}$$

$$\therefore 3x^{\frac{2}{3}}y^{\frac{1}{3}} + 3x^{\frac{1}{3}}y^{\frac{2}{3}} = 144,$$

$$\text{and } x^{\frac{1}{3}}y^{\frac{1}{3}}(x^{\frac{1}{3}} + y^{\frac{1}{3}}) = 48;$$

$$\therefore x^{\frac{1}{3}}y^{\frac{1}{3}} = \frac{48}{x^{\frac{1}{3}}+y^{\frac{1}{3}}} = 8,$$

$$\text{and } xy = 512.$$

$$\text{Now } x^3 + 2xy + y^3 = 5184;$$

$$\text{and } 4xy = 2048;$$

$$\therefore x^3 - 2xy + y^3 = 3136;$$

$$\text{and } \therefore x - y = \pm 56,$$

$$\text{but } x + y = 72;$$

$$\therefore x = 64 \text{ or } 8.$$

$$\text{and } y = 8 \text{ or } 64.$$

$$21. \quad \left. \begin{array}{l} \text{Given } 4x^2 + \frac{5}{2} = \frac{x^2}{y} + 10y \\ \text{and } x^2 + 3y = 55 \end{array} \right\} \begin{array}{l} \text{to find the values of} \\ x \text{ and } y. \end{array}$$

$$\text{by transposition } 4x^2 - \frac{x^2}{y} = 10y - \frac{5}{2};$$

$$\therefore \frac{4x^2y - x^2}{y} = \frac{20y - 5}{2};$$

$$\text{or } \frac{x^2}{y} (4y - 1) = \frac{5}{2} (4y - 1);$$

$$\therefore \frac{x^2}{y} = \frac{5}{2}, \quad \text{unless } y = \frac{1}{4}$$

$$\text{and } x^2 = \frac{5y}{2}.$$

Substituting this value in the second equation,

$$\frac{5y}{2} + 3y = 55,$$

$$11y = 110;$$

$$\text{and } y = 10;$$

$$\therefore x^2 = \frac{5y}{2} = \frac{50}{2} = 25,$$

$$\text{and } x = \pm 5.$$

22. Given $\frac{1}{x + \sqrt{2-x^2}} + \frac{1}{x - \sqrt{2-x^2}} = ax$, to find the values of x .

$$\text{then } \frac{x - \sqrt{2-x^2} + x + \sqrt{2-x^2}}{2x^2 - 2} = ax;$$

$$\therefore \frac{2x}{2x^2 - 2} = ax;$$

$$\frac{1}{x^2 - 1} = a,$$

$$\text{and } x^2 - 1 = \frac{1}{a},$$

$$x^2 = \frac{1}{a} + 1 = \frac{a+1}{a};$$

$$\therefore x = \pm \sqrt{\frac{a+1}{a}}.$$

23. Given $\frac{x}{\sqrt{a^2+x^2}-x} = b$, to find the values of x .

$$\text{then } x : \sqrt{a^2+x^2} - x :: b : 1,$$

$$x : \sqrt{a^2+x^2} :: b : b+1. \quad (\text{Alg. 179.})$$

$$x^2 : a^2 + x^2 :: b^2 : b^2 + 2b + 1;$$

$$x^2 : a^2 :: b^2 : 2b + 1;$$

$$\therefore x^2 = \frac{a^2 b^2}{2b + 1},$$

$$\text{and } x = \pm \frac{ab}{\sqrt{2b + 1}}.$$

24. Given $\sqrt{y} - \sqrt{a-x} = \sqrt{y-x}$ } to find the values
and $\sqrt{y-x} + \sqrt{a-x} : \sqrt{a-x} :: 5 : 2$ } of x and y .

$$\text{From the first equation, } \sqrt{y} = \sqrt{y-x} + \sqrt{a-x};$$

Substituting this value in the proportion,

$$\therefore \sqrt{y} : \sqrt{a-x} :: 5 : 2;$$

$$\text{but } \sqrt{a-x} : \sqrt{y-x} :: 2 : 3;$$

$$\therefore \sqrt{y} : \sqrt{y-x} :: 5 : 3,$$

$$\text{and } y : y-x :: 25 : 9. \quad (\text{Alg. 188}).$$

$$y : x :: 25 : 16. \quad (\text{Alg. 181.});$$

$$\therefore y = \frac{25x}{16},$$

$$\text{by substitution, } \sqrt{\frac{25x}{16}} - \sqrt{a-x} = \sqrt{\frac{9x}{16}},$$

$$\text{or } \frac{5\sqrt{x} - 3\sqrt{x}}{4} = \sqrt{a-x};$$

$$\therefore \frac{\sqrt{x}}{2} = \sqrt{a-x}.$$

Squaring both sides,

$$\frac{x}{4} = a - x,$$

$$\text{and } x = 4a - 4x;$$

$$\therefore 5x = 4a,$$

$$\text{and } x = \frac{4a}{5};$$

$$\therefore y = \frac{25x}{16} = \frac{100a}{80} = \frac{5a}{4}.$$

$$\left. \begin{array}{l} 25. \text{ Given } x^{\frac{4}{3}} + y^{\frac{2}{3}} = 20, \\ \text{and } x^{\frac{2}{3}} + y^{\frac{1}{3}} = 6, \end{array} \right\} \text{to find the values of } x \text{ and } y.$$

Squaring the second equation, we have,

$$x^{\frac{4}{3}} + 2x^{\frac{2}{3}}y^{\frac{1}{3}} + y^{\frac{2}{3}} = 36,$$

$$x^{\frac{4}{3}} + y^{\frac{2}{3}} = 20.$$

$$\therefore 2x^{\frac{2}{3}}y^{\frac{1}{3}} = 16,$$

and $x^{\frac{2}{3}} - 2x^{\frac{2}{3}}y^{\frac{1}{3}} + y^{\frac{2}{3}} = 4$; and, extracting the square root on both sides,

$$\begin{aligned} x^{\frac{2}{3}} - y^{\frac{2}{3}} &= \pm 2, \\ \text{but } x^{\frac{2}{3}} + y^{\frac{2}{3}} &= 6; \\ \hline \therefore x^{\frac{2}{3}} &= 4 \text{ or } 2, \\ x &= \pm 8 \text{ or } \pm \sqrt[3]{8}, \\ y^{\frac{2}{3}} &= 2 \text{ or } 4, \\ \text{and } y &= 32 \text{ or } 1024. \end{aligned}$$

26. Given $x^4 + 2x^2y^2 + y^4 = 1296 - 4xy(x^2 + xy + y^2)$
and $x - y = 4$,
to find the values of x and y .

$$\begin{aligned} x^4 + 2x^2y^2 + y^4 &= 1296 - 4x^2y + 4x^2y^2 - 4xy^2, \\ \therefore x^4 + 4x^2y + 6x^2y^2 + 4xy^2 + y^4 &= 1296; \end{aligned}$$

and extracting the fourth root,

$$\begin{aligned} x + y &= \pm 6, \\ \text{but } x - y &= 4; \\ \hline \therefore x &= 5 \text{ or } -1, \\ \text{and } y &= 1 \text{ or } -5. \end{aligned}$$

27. Given $\frac{\sqrt{a} - \sqrt{a-x}}{\sqrt{a} + \sqrt{a-x}} = a$, to find the value of x .

1. Multiplying numerator and denominator by $(\sqrt{a} - \sqrt{a-x})$

$$\frac{(\sqrt{a} - \sqrt{a-x})^2}{x} = a,$$

$$(\sqrt{a} - \sqrt{a-x})^2 = ax;$$

$$\text{and } \sqrt{a} - \sqrt{a-x} = \sqrt{ax};$$

$$\therefore \sqrt{a} - \sqrt{ax} = \sqrt{a-x}.$$

Squaring both sides,

$$a - 2\sqrt{a^2x} + ax = a - x;$$

$$ax + x = 2 \sqrt{a^2 x};$$

$$\therefore \sqrt{x} (a + 1) = 2a \text{ (dividing both sides by } \sqrt{x} \text{),}$$

$$x = \frac{4a^2}{(a + 1)^2}.$$

2. Thus, by proportion,

$$\sqrt{a} - \sqrt{a-x} : \sqrt{a} + \sqrt{a-x} :: a : 1,$$

$$\sqrt{a} : \sqrt{a-x} : 1 + a :: 1 - a. \text{ (Alg. 182.)}$$

$$a : a - x :: 1 + 2a + a^2 : 1 - 2a + a^2;$$

$$a : x :: 1 + 2a + a^2 : 4a,$$

$$x = \frac{4a^2}{(a+1)^2}.$$

$$28. \text{ Given } \frac{\sqrt{x} + \sqrt{x-y}}{\sqrt{x} + \sqrt{x-y}} = 4, \left. \begin{array}{l} \text{to find the values of} \\ x \text{ and } y. \end{array} \right\}$$

$$\text{and } \sqrt{x} : \sqrt{y} :: \sqrt{y} : 4$$

$$\sqrt{x} + \sqrt{x-y} : \sqrt{x} - \sqrt{x-y} :: 4 : 1,$$

$$\sqrt{x} : \sqrt{x-y} :: 5 : 3. \text{ (Alg. 182.)}$$

$$\therefore x : x - y :: 25 : 9,$$

$$\text{and } x : y :: 25 : 16;$$

$$\text{but } y : x :: 16 : y;$$

$$\therefore y = 25,$$

$$\text{and } x = \frac{y^2}{16} = \frac{625}{16}.$$

$$29. \text{ Given } \frac{1}{y} - \frac{1}{x} = \frac{1}{4} \left. \begin{array}{l} \text{to find the values of } x \text{ and } y. \\ \text{and } x^2 y - xy^2 = 16 \end{array} \right\}$$

$$x - y = \frac{xy}{4} \text{ (from the first equation,)}$$

$$\text{and } x - y = \frac{16}{xy} \text{ (from the second equation;)}$$

$$\therefore \frac{xy}{4} = \frac{16}{xy}$$

$$x^2 y^2 = 64,$$

$$\text{and } xy = 8.$$

By substitution in the first equation,

$$x - y = 2,$$

$$x^2 - 2xy + y^2 = 4,$$

$$4xy = 32.$$

$$\hline x^2 + 2xy + y^2 = 36$$

$$x + y = \pm 6$$

$$x - y = 2.$$

$$\hline x = 4 \text{ or } -2$$

$$y = 2 \text{ or } -4.$$

30. Given $\frac{\sqrt{4x+1} + \sqrt{4x}}{\sqrt{4x+1} - \sqrt{4x}} = 9$, to find the value of x .

$$\sqrt{4x+1} + \sqrt{4x} : \sqrt{4x+1} - \sqrt{4x} :: 9 : 1,$$

$$\sqrt{4x+1} : \sqrt{4x} :: 5 : 4. \quad (\text{Alg. 182.})$$

$$4x + 1 : 4x :: 25 : 16;$$

$$4x : 1 :: 16 : 9;$$

$$\therefore 4x = \frac{16}{9},$$

$$\text{and } x = \frac{4}{9}.$$

31. Given $\frac{a+x+\sqrt{2ax+x^2}}{a+x-\sqrt{2ax+x^2}} = b$, to find the values of x .

$$a+x+\sqrt{2ax+x^2} : a+x-\sqrt{2ax+x^2} :: b : 1,$$

$$a+x : \sqrt{2ax+x^2} :: b+1 : b-1. \quad (\text{Alg. 182.})$$

$$a^2 + 2ax + x^2 : 2ax + x^2 :: b^2 + 2b + 1 : b^2 - 2b + 1$$

$$a^2 + 2ax + x^2 : a^2 :: b^2 + 2b + 1 : 4b.$$

$$\begin{aligned}
 a + x : a &:: b + 1 : \pm 2\sqrt{b}, \\
 x : a &:: b \mp 2\sqrt{b} + 1 : \pm 2\sqrt{b}; \\
 \therefore x &= \pm a \cdot \frac{(\sqrt{b} \mp 1)^2}{2\sqrt{b}}.
 \end{aligned}$$

32. Given, $xy^3 - x^3y = 216$ } to find the values of x
 and $x^2y - xy^2 = 6$ } and y .

Dividing the first equation by the second,

$$\text{we have } x^2y^2 = 36;$$

$$\text{and } xy = 6.$$

$$\text{from the second equation } x - y = \frac{6}{xy};$$

$$\therefore x - y = 1.$$

Squaring both sides,

$$x^2 - 2xy + y^2 = 1,$$

$$4xy = 24;$$

$$\therefore x^2 + 2xy + y^2 = 25,$$

$$\text{and } x + y = \pm 5;$$

$$\text{but } x - y = 1;$$

$$\therefore x = 3 \text{ or } -2,$$

$$\text{and } y = 2 \text{ or } -3.$$

33. Given $x^3 + x\sqrt[3]{xy^3} = 208$ } to find the values of
 and $y^3 + y\sqrt[3]{x^3y} = 1053$ } x and y .

$$x^{\frac{4}{3}} \cdot (x^{\frac{2}{3}} + y^{\frac{2}{3}}) = 208,$$

$$y^{\frac{4}{3}} \cdot (x^{\frac{2}{3}} + y^{\frac{2}{3}}) = 1053.$$

Dividing the former by the latter equation,

$$\frac{x^{\frac{4}{3}}}{y^{\frac{4}{3}}} = \frac{208}{1053} = \frac{16}{81},$$

$$\frac{x^{\frac{2}{3}}}{y^{\frac{2}{3}}} = \frac{4}{9}; \text{ and } y^{\frac{2}{3}} = \frac{9x^{\frac{2}{3}}}{4}.$$

Substituting this value of $y^{\frac{2}{3}}$ in the first equation,

$$x^3 + \frac{x^{\frac{4}{3}} \times 9x^{\frac{2}{3}}}{4} = 208;$$

$$\text{or } \frac{13x^3}{4} = 208;$$

$$\therefore x^3 = 4 \times 16 = 64,$$

$$\text{and } x = \pm 4;$$

$$\therefore y^{\frac{2}{3}} = \frac{9x^{\frac{2}{3}}}{4},$$

$$\text{and } y = \frac{27x}{8} = \pm 27.$$

34. Given $x^{\frac{3}{2}} + x^{\frac{1}{2}}y^{\frac{3}{2}} + y^{\frac{3}{2}} = 1009$ } to find the
 and $x^3 + x^{\frac{3}{2}}y^{\frac{3}{2}} + y^3 = 582193$ }
 values of x and y .

Dividing the second equation by the first,

$$\text{we have } x^{\frac{3}{2}} - x^{\frac{1}{2}}y^{\frac{3}{2}} + y^{\frac{3}{2}} = 577 \text{ (c),}$$

$$\text{but } x^{\frac{3}{2}} + x^{\frac{1}{2}}y^{\frac{3}{2}} + y^{\frac{3}{2}} = 1009 \text{ (b);}$$

$$\therefore 2x^{\frac{1}{2}}y^{\frac{3}{2}} = 432,$$

$$\text{and } x^{\frac{1}{2}}y^{\frac{3}{2}} = 216 \text{ (a),}$$

adding (a) to (b) we have

$$x^{\frac{3}{2}} + 2x^{\frac{1}{2}}y^{\frac{3}{2}} + y^{\frac{3}{2}} = 1225$$

$$\text{or } x^{\frac{3}{2}} + y^{\frac{3}{2}} = \pm 35.$$

Subtracting (a) from (c) we have

$$x^{\frac{3}{2}} - 2x^{\frac{1}{2}}y^{\frac{3}{2}} + y^{\frac{3}{2}} = 361,$$

$$\text{or } x^{\frac{3}{2}} - y^{\frac{3}{2}} = \pm 19,$$

$$\text{hence } x^{\frac{3}{2}} + y^{\frac{3}{2}} = \pm 35;$$

$$\text{and } x^{\frac{3}{2}} - y^{\frac{3}{2}} = \pm 19;$$

$$\therefore 2x^{\frac{3}{2}} = 54 \text{ or } 16,$$

$$x^{\frac{3}{2}} = 27 \text{ or } 8;$$

$$\text{and } x = 81 \text{ or } 16,$$

$$\text{and } y = 16 \text{ or } 81.$$

35. Given $x^3 + y^3 + xy(x + y) = 68$
and $x^3 + y^3 - 3x^2 = 12 + 3y^2$ } to find the values
of x and y .

Multiplying the first equation by 3,

$$3x^3 + 3y^3 + 3x^2y + 3xy^2 = 204,$$

$$\text{and } x^3 + y^3 - 3x^2 - 3y^2 = 12.$$

$$\text{then } x^3 + 3x^2y + 3xy^2 + y^3 = 216,$$

$$x + y = 6.$$

Substituting this value for $x + y$ in the first equation;

$$\therefore x^3 + 6xy + y^3 = 68,$$

$$\text{but } x^3 + 2xy + y^3 = 36.$$

$$4xy = 32,$$

$$\text{then } x^3 - 2xy + y^3 = 4,$$

$$\text{and } x - y = \pm 2,$$

$$\text{but } x + y = 6;$$

$$\therefore x = 4 \text{ or } 2,$$

$$\text{and } y = 2 \text{ or } 4.$$

36. Given $xy \cdot \overline{x + y} = 84$, } to find the values of x
and $x^2y^2 \cdot \overline{x^2 + y^2} = 3600$, } and y .

From the first equation $x^2y + xy^2 = 84$;

$$\therefore x^4y^3 + 2x^3y^2 + x^2y^4 = 7056.$$

From the second equation $x^4y^3 + x^2y^4 = 3600$;

$$\therefore 2x^3y^3 = 3456,$$

$$x^2y^3 = 1728,$$

$$\text{and } xy = 12.$$

$$\text{Now } x + y = \frac{84}{xy} = 7;$$

$$\therefore x^2 + 2xy + y^2 = 49,$$

$$\quad \quad \quad 4xy = 48;$$

$$\therefore x^2 - 2xy + y^2 = 1,$$

$$\text{and } x - y = \pm 1,$$

$$\text{but } x + y = 7;$$

$$\therefore x = 4 \text{ or } 3,$$

$$\text{and } y = 3 \text{ or } 4.$$

$$37. \text{ Given } \left. \begin{array}{l} \frac{x^2 + xy + y^2}{x + y} = 7 \text{ (a),} \\ \text{and } \frac{x^2 - xy + y^2}{x - y} = 9 \text{ (b),} \end{array} \right\} \begin{array}{l} \text{to find the values of } x \\ \text{and } y. \end{array}$$

Dividing (a) by (b) we have,

$$\frac{x^2 - y^2}{x^2 + y^2} = \frac{7}{9};$$

$$\therefore x^2 + y^2 : x^2 - y^2 :: 9 : 7$$

$$x^2 : y^2 :: 16 : 2 \quad (\text{Alg. 162.})$$

$$x^2 : y^2 :: 8 : 1,$$

$$x : y :: 2 : 1,$$

$$\text{and } x = 2y;$$

$$\therefore \frac{4y^2 + 2y^2 + y^2}{3y} = 7,$$

$$\text{and } \frac{7y}{3} = 7;$$

$$\therefore y = 3,$$

$$\text{and } x = 2y = 6.$$

Solutions of Affected Quadratics, involving only one unknown Quantity.

36. Given $\frac{\sqrt{x} + 9}{\sqrt{x}} = \frac{\sqrt{9x} - 3\frac{1}{2}}{9 - \sqrt{x}}$, to find the values of x .

$$\text{then } 81 - x = 3x - \frac{19\sqrt{x}}{5},$$

$$\text{or } 4x - \frac{19\sqrt{x}}{5} = 81;$$

$$\therefore x - \frac{19\sqrt{x}}{20} + \frac{361}{1600} = \frac{81}{4} + \frac{361}{1600} = \frac{32400 + 361}{1600} = \frac{32761}{1600};$$

$$\therefore \sqrt{x} - \frac{19}{40} = \pm \frac{181}{40},$$

$$\sqrt{x} = \frac{200}{40} \text{ or } -\frac{162}{40},$$

$$\sqrt{x} = 5 \text{ or } -\frac{81}{20};$$

$$\therefore x = 25 \text{ or } \frac{6561}{400}.$$

37. Given $\frac{x + \sqrt{x}}{x - \sqrt{x}} = \frac{x^2 - x}{4}$, to find the values of x .

Dividing both sides by $x + \sqrt{x}$,

$$\frac{1}{x - \sqrt{x}} = \frac{x - \sqrt{x}}{4};$$

$$\therefore (x - \sqrt{x})^2 = 4,$$

$$\text{and } x - \sqrt{x} = \pm 2.$$

Completing the square,

$$x - \sqrt{x} + \frac{1}{4} = \frac{1}{4} \pm 2 = \frac{9}{4} \text{ or } -\frac{7}{4};$$

$$\therefore \sqrt{x} - \frac{1}{2} = \pm \frac{3}{2} \text{ or } \pm \frac{\sqrt{-7}}{2},$$

$$\text{and } \sqrt{x} = 2 \text{ or } -1 \text{ or } \frac{1 \pm \sqrt{-7}}{2}.$$

Squaring both sides,

$$x = 4 \text{ or } 1 \text{ or } \frac{1 \pm 2\sqrt{-7} - 7}{4},$$

$$= 4 \text{ or } 1 \text{ or } \frac{-6 \pm 2\sqrt{-7}}{4},$$

$$= 4 \text{ or } 1 \text{ or } \frac{-3 \pm \sqrt{-7}}{2}.$$

38. Given $\frac{x - \sqrt{x+1}}{x + \sqrt{x+1}} = \frac{5}{11}$, to find the values of x .

$$x + \sqrt{x+1} : x - \sqrt{x+1} :: 11 : 5,$$

$$\text{then } x : \sqrt{x+1} :: 8 : 3;$$

$$\therefore x^2 : x+1 :: 64 : 9,$$

$$\text{and } x^2 = \frac{64}{9}x + \frac{64}{9};$$

$$\therefore x^2 - \frac{64x}{9} + \frac{32}{9} = \frac{1024}{81} + \frac{64}{9} = \frac{1600}{81},$$

$$\text{then } x - \frac{32}{9} = \pm \frac{40}{9};$$

$$\text{and } \therefore x = 8 \text{ or } -\frac{8}{9}$$

39. Given $5 \cdot \frac{3x-1}{1+5\sqrt{x}} + \frac{2}{\sqrt{x}} = 3\sqrt{x}$, to find the values of x .

$$\text{Now } 15x - 5 + \frac{2 + 10\sqrt{x}}{\sqrt{x}} = 3\sqrt{x} + 15x;$$

$$\therefore 3x + 5\sqrt{x} = 10\sqrt{x} + 2,$$

$$\text{and } 3x - 5\sqrt{x} = 2,$$

then $x - \frac{5\sqrt{x}}{3} = \frac{2}{3}$, and completing the square,

$$x - \frac{5\sqrt{x}}{3} + \frac{25}{36} = \frac{24 + 25}{36} = \frac{49}{36};$$

$$\therefore \sqrt{x} - \frac{5}{6} = \pm \frac{7}{6},$$

$$\sqrt{x} = 2 \text{ or } -\frac{1}{3},$$

$$\text{and } x = 4 \text{ or } \frac{1}{9}.$$

40. Given $\sqrt{x^5} - \frac{40}{\sqrt{x}} = 3x$, to find the values of x .

Clearing of fractions,

$$x^3 - 40 = 3x^{\frac{3}{2}},$$

then $x^3 - 3x^{\frac{3}{2}} = 40$; and completing the square,

$$x^3 - 3x^{\frac{3}{2}} + \left(\frac{3}{2}\right)^2 = \frac{9}{4} + 40 = \frac{169}{4};$$

$$\therefore x^{\frac{3}{2}} - \frac{3}{2} = \pm \frac{13}{2},$$

$$x^{\frac{3}{2}} = 8 \text{ or } -5,$$

$$\text{and } x = 4 \text{ or } \sqrt[3]{-5}^{\frac{2}{3}}.$$

41. Given $x^{\frac{4}{3}} + 7x^{\frac{2}{3}} = 44$, to find the values of x .

$$\text{Here } x^{\frac{4}{3}} + 7x^{\frac{2}{3}} + \frac{49}{4} = 44 + \frac{49}{4} = \frac{225}{4}$$

$$\text{then } x^{\frac{2}{3}} + \frac{7}{2} = \pm \frac{15}{2},$$

$$\text{and } x^{\frac{2}{3}} = 4 \text{ or } -11;$$

$$\therefore x = \pm 8 \text{ or } \sqrt[3]{-11}^{\frac{3}{2}}.$$

42. Given $4x^{\frac{1}{3}} + x^{\frac{1}{6}} = 39$, to find the values of x .

$$\text{Dividing by 4, } x^{\frac{1}{3}} + \frac{x^{\frac{1}{6}}}{4} = \frac{39}{4},$$

$$\text{then } x^{\frac{1}{3}} + \frac{x^{\frac{1}{6}}}{4} + \frac{1}{64} = \frac{625}{64},$$

$$\text{and } x^{\frac{1}{6}} + \frac{1}{8} = \pm \frac{25}{8};$$

$$\therefore x^{\frac{1}{6}} = 3 \text{ or } -\frac{13}{4},$$

$$\text{and } x = 729 \text{ or } \sqrt[6]{-\frac{13}{4}}.$$

43. Given $3x^6 + 42x^3 = 3321$, to find the values of x .

$$\text{Now } x^6 + 14x^3 = 1107;$$

$$\therefore x^6 + 14x^3 + 49 = 1107 + 49 = 1156,$$

$$\text{and } x^3 + 7 = \pm 34;$$

$$\therefore x^3 = 27 \text{ or } -41,$$

$$\text{and } x = 3 \text{ or } -\sqrt[3]{41}.$$

44. Given $\frac{8}{x^3} + 2 = \frac{17}{x^{\frac{3}{2}}}$, to find the values of x .

Dividing by 8, and transposing

$$\frac{1}{x^3} - \frac{17}{8x^{\frac{3}{2}}} = -\frac{1}{4},$$

$$\frac{1}{x^3} - \frac{17}{8x^{\frac{3}{2}}} + \frac{289}{256} = \frac{289}{256} - \frac{1}{4} = \frac{225}{256},$$

$$\frac{1}{x^{\frac{3}{2}}} - \frac{17}{16} = \pm \frac{15}{16},$$

$$\frac{1}{x^{\frac{3}{2}}} = 2 \text{ or } \frac{1}{8};$$

$$\therefore x^{\frac{3}{2}} = 8 \text{ or } \frac{1}{2},$$

$$x = 4 \text{ or } \sqrt[4]{\frac{1}{4}}.$$

45. Given $x^{\frac{1}{3}} + \frac{41\sqrt[3]{-x}}{x} = \frac{97}{\sqrt[3]{x^2}} + x^{\frac{5}{6}}$, to find the values of x .

$$\text{or } x^{\frac{1}{3}} + \frac{41}{x^{\frac{2}{3}}} = \frac{97}{x^{\frac{2}{3}}} + x^{\frac{5}{6}},$$

$$\text{or } x^{\frac{1}{3}} + 41 = 97 + x^{\frac{5}{6}},$$

$$\therefore x^{\frac{1}{3}} - x^{\frac{5}{6}} = 56,$$

$$x^{\frac{1}{3}} - x^{\frac{5}{6}} + \frac{1}{4} = \frac{224 + 1}{4} = \frac{225}{4};$$

$$\therefore x^{\frac{1}{3}} - \frac{1}{2} = \pm \frac{15}{2},$$

$$\text{and } x^{\frac{1}{3}} = 8 \text{ or } -7;$$

$$\therefore x = 4 \text{ or } -7\frac{2}{3}.$$

46. Given $\sqrt{\frac{1}{x^4}} + \sqrt{\frac{1}{x}} = \frac{3 - \sqrt[3]{x^2}}{x}$, to find the values of x .

$$\frac{1}{x^{\frac{3}{2}}} + \frac{1}{x^{\frac{1}{2}}} = \frac{3 - x^{\frac{2}{3}}}{x},$$

$$\text{then } x^{\frac{1}{3}} + x^{\frac{2}{3}} = 3 - x^{\frac{2}{3}},$$

$$2x^{\frac{2}{3}} + x^{\frac{1}{3}} = 3,$$

$$x^{\frac{2}{3}} + \frac{x^{\frac{1}{3}}}{2} + \frac{1}{16} = \frac{24 + 1}{16} = \frac{25}{16},$$

$$x^{\frac{1}{3}} + \frac{1}{4} = \pm \frac{5}{4},$$

$$x^{\frac{1}{3}} = 1 \text{ or } -\frac{3}{2};$$

$$\therefore x = 1 \text{ or } -\frac{27}{8}.$$

47. Given $3x^{\frac{2}{3}} - 4x^{\frac{1}{3}} = 4$, to find the values of x .

$$3x^{\frac{2}{3}} - 4x^{\frac{1}{3}} = 4,$$

$$x^{\frac{2}{3}} - \frac{4x^{\frac{1}{3}}}{3} = \frac{4}{3},$$

$$x^{\frac{2}{3}} - \frac{4x^{\frac{1}{3}}}{3} + \frac{4}{9} = \frac{12 + 4}{9} = \frac{16}{9};$$

$$x^{\frac{2}{3}} - \frac{2}{3} = \pm \frac{4}{3},$$

$$x^{\frac{2}{3}} = 2 \text{ or } -\frac{2}{3},$$

$$x^{\frac{2}{3}} = 8 \text{ or } -\frac{8}{27};$$

$$\text{and } x = 8^{\frac{3}{2}} \text{ or } -\frac{8}{27}^{\frac{3}{2}}.$$

48. Given $adx - acx^2 = bcx - bd$, to find the values of x .

$$acx^2 + bcx - adx = bd$$

$$x^2 + \frac{bc - ad}{ac} x = \frac{bd}{ac}$$

$$\begin{aligned} x^2 + \frac{bc - ad}{ac} x + \left(\frac{bc - ad}{2ac} \right)^2 &= \frac{bd}{ac} + \frac{b^2c^2 + 2abcd + a^2d^2}{4a^2c^2} \\ &= \frac{4abcd + b^2c^2 - 2abcd + a^2d^2}{4a^2c^2} \\ &= \frac{b^2c^2 + 2abcd + a^2d^2}{4a^2c^2}; \end{aligned}$$

$$\therefore x + \frac{bc - ad}{2ac} = \pm \frac{bc + ad}{2ac};$$

$$\text{and } x = \frac{2ad}{2ac} \text{ or } -\frac{2bc}{2ac}$$

$$= \frac{d}{c} \text{ or } -\frac{b}{a}.$$

49. Given $\frac{a^2x^2}{b^2} - \frac{2ax}{c} + \frac{d^2}{c^2} = 0$, to find the values of x .

$$a^2x^2 - \frac{2ab^2x}{c} = -\frac{b^2d^2}{c^2},$$

$$a^2x^2 - \frac{2ab^2x}{c} + \frac{b^4}{c^2} = \frac{b^4}{c^2} - \frac{b^2d^2}{c^2};$$

$$\therefore ax - \frac{b^2}{c} = \frac{\pm \sqrt{b^4 - b^2d^2}}{c} = \frac{\pm b \sqrt{b^2 - d^2}}{c}$$

$$ax = \frac{b^2 \pm b \sqrt{b^2 - d^2}}{c} = b \cdot \frac{b \pm \sqrt{b^2 - d^2}}{c}$$

$$\therefore x = \frac{b}{a} \cdot \frac{b \pm \sqrt{b^2 - d^2}}{c}.$$

50. Given, $9a^4b^4x^2 - 6a^3b^3x = b^2$ to find the values of x .

$$x^2 - \frac{2x}{3ab^2} = \frac{1}{9a^4b^2},$$

$$x^2 - \frac{2x}{3ab^2} + \frac{1}{9a^2b^4} = \frac{1}{9a^2b^4} + \frac{1}{9a^4b^2}$$

$$= \frac{a^2 + b^2}{9a^4b^4};$$

$$x - \frac{1}{3ab^2} = \frac{\pm \sqrt{a^2 + b^2}}{3a^2b^2},$$

$$x = \frac{a \pm \sqrt{a^2 + b^2}}{3a^2b^2}.$$

51. Given, $\overline{a+b} \cdot x^2 = cx + \frac{ac}{a+b}$, to find the values of x .

$$\overline{a+b} \cdot x^2 - cx = \frac{ac}{a+b};$$

$$x^2 - \frac{cx}{a+b} = \frac{ac}{(a+b)^2};$$

$$x^2 - \frac{cx}{a+b} + \frac{c^2}{4(a+b)^2} = \frac{ac}{a+b} + \frac{c^2}{4(a+b)^2} = \frac{c^2 + 4ac}{4(a+b)^2}$$

$$x - \frac{c}{2(a+b)} = \pm \frac{\sqrt{c^2 + 4ac}}{2(a+b)};$$

$$\text{and } x = \frac{c \pm \sqrt{c^2 + 4ac}}{2(a+b)}.$$

The succeeding five equations merely require attention to the ordinary rules, as laid down by Dr. Bland; they are long and tedious, without a single point worthy of notice, and would consequently occupy room to no purpose.

57. Given $x + \sqrt{x} : x - \sqrt{x} :: 3\sqrt{x} + 6 : 2\sqrt{x}$
to find the values of x .

$$\text{then } \sqrt{x} + 1 : \sqrt{x} - 1 :: 3\sqrt{x} + 6 : 2\sqrt{x};$$

$$\text{or } 2x + 2\sqrt{x} = 3x + 3\sqrt{x} - 6,$$

$$x + \sqrt{x} = 6;$$

$$x + \sqrt{x} + \frac{1}{4} = \frac{25}{4}$$

$$\sqrt{x} + \frac{1}{2} = \pm \frac{5}{2}.$$

$$\sqrt{x} = 2 \text{ or } 3;$$

$$\therefore x = 4 \text{ or } 9.$$

58. Given $x^2 + 11 + \sqrt{x^2 + 11} = 42$, to find the values of x .

$$x^2 + 11 + \sqrt{x^2 + 11} + \frac{1}{4} = \frac{169}{4},$$

$$\sqrt{x^2 + 11} + \frac{1}{2} = \pm \frac{13}{2}$$

$$\sqrt{x^2 + 11} = 6 \text{ or } -7,$$

$$x^2 + 11 = 36 \text{ or } 49,$$

$$x^2 = 25 \text{ or } 38;$$

$$\therefore x = \pm 5 \text{ or } \pm \sqrt{38}$$

59. Given, $\sqrt{x-5}^3 - 3 \cdot \sqrt{x-5}^{\frac{3}{2}} = 40$, to find the values of x .

$$\sqrt{x-5}^3 - 3 \cdot \sqrt{x-5}^{\frac{3}{2}} + \frac{9}{4} = 40 + \frac{9}{4} = \frac{169}{4}$$

$$\sqrt{x-5}^{\frac{3}{2}} - \frac{3}{2} = \pm \frac{13}{2}$$

$$\sqrt{x-5}^{\frac{3}{2}} = 8 \text{ or } -5$$

$$x - 5 = 4 \text{ or } \sqrt{-5}^{\frac{2}{3}}$$

$$x = 9 \text{ or } \sqrt{-5}^{\frac{2}{3}} + 5.$$

60. Given $x + \sqrt{x+6} = 2 + 3\sqrt{x+6}$, to find the values of x .

$$x - 2\sqrt{x+6} = 2$$

$$x + 6 - 2\sqrt{x+6} = 8$$

$$x + 6 - 2\sqrt{x+6} + 1 = 9$$

$$\sqrt{x+6} - 1 = \pm 3$$

$$\sqrt{x+6} = 4 \text{ or } -2$$

$$x + 6 = 16 \text{ or } 4.$$

$$\text{and } x = 10 \text{ or } -2.$$

61. Given $\sqrt{x^2+5}^2 - 4x^2 = 160$, to find the values of x .

$$\text{then } \sqrt{x^2+5}^2 - 4x^2 - 20 = 140$$

$$\sqrt{x^2+5}^2 - 4(x^2+5) + 4 = 144$$

$$x^2 + 5 - 2 = \pm 12$$

$$x^2 + 5 = 14 \text{ or } -10$$

$$x^2 = 9 \text{ or } -15$$

$$\text{and } x = \pm 3 \pm \sqrt{-15}.$$

62. Given $x^2 - 7x + \sqrt{x^2 - 7x + 18} = 24$, to find the values of x .

$$x^2 - 7x + 18 + \sqrt{x^2 - 7x + 18} + \frac{1}{4} = 42 + \frac{1}{4} = \frac{169}{4}$$

$$\therefore \sqrt{x^2 - 7x + 18} + \frac{1}{2} = \pm \frac{13}{2}$$

$$\sqrt{x^2 - 7x + 18} = 6 \text{ or } -7,$$

$$\text{and } x^2 - 7x + 18 = 36 \text{ or } 49.$$

$$\therefore x^2 - 7x = 18 \text{ or } 31,$$

$$x^2 - 7x + \frac{49}{4} = \frac{121}{4} \text{ or } \frac{173}{4}$$

$$x - \frac{7}{2} = \pm \frac{11}{2} \text{ or } \pm \frac{\sqrt{173}}{2};$$

$$\therefore x = 9 \text{ or } 2 \text{ or } \frac{7 \pm \sqrt{173}}{2}.$$

63. Given $4x^2 - 9x + \sqrt{4x^2 - 9x + 11} = 5$, to find the values of x .

$$\text{Now } 4x^2 - 9x - \sqrt{4x^2 - 9x + 11} = -5,$$

$$4x^2 - 9x + 11 - \sqrt{4x^2 - 9x + 11} + \frac{1}{4} = 6 + \frac{1}{4} = \frac{25}{4}$$

$$\therefore \sqrt{4x^2 - 9x + 11} - \frac{1}{2} = \pm \frac{5}{2}$$

$$\sqrt{4x^2 - 9x + 11} = 3 \text{ or } -2$$

$$\text{and } 4x^2 - 9x + 11 = 9 \text{ or } 4,$$

$$\therefore 4x^2 - 9x = -2 \text{ or } -7$$

$$x^2 - \frac{9x}{4} = -\frac{2}{4} \text{ or } -\frac{7}{4}$$

$$x^2 - \frac{9x}{4} + \frac{81}{64} = \frac{49}{64} \text{ or } -\frac{31}{64}$$

$$x - \frac{9}{8} = \pm \frac{7}{8} \text{ or } \pm \frac{\sqrt{-31}}{8}$$

$$\therefore x = 2 \text{ or } \frac{1}{4} \text{ or } \frac{9 \pm \sqrt{-31}}{8}.$$

64. Given $x^2 + \sqrt{5x+x^2} = 42 - 5x$, to find the values of x .

$$\text{Here } 5x + x^2 + \sqrt{5x+x^2} = 42$$

$$5x + x^2 + \sqrt{5x+x^2} + \frac{1}{4} = \frac{169}{4}$$

$$\sqrt{5x+x^2} + \frac{1}{2} = \pm \frac{13}{2}$$

$$\sqrt{5x+x^2} = 6 \text{ or } -7,$$

$$\therefore 5x + x^2 = 36 \text{ or } 49$$

$$\text{Again, } x^2 + 5x + \frac{25}{4} = 36 + \frac{25}{4} \text{ or } 49 + \frac{25}{4}$$

$$= \frac{169}{4} \text{ or } \frac{221}{4}$$

$$x + \frac{5}{2} = \pm \frac{13}{2} \text{ or } \pm \frac{\sqrt{221}}{2}$$

$$= 4 \text{ or } -9 \text{ or } \frac{-5 \pm \sqrt{221}}{2}$$

65. Given $\frac{2}{(x+2)^{\frac{3}{2}}} + \frac{\sqrt{x+2}}{2} = \frac{17}{4\sqrt{x+2}}$, to find the values of x .

$$\frac{1}{(x+2)^{\frac{3}{2}}} + \frac{1}{4} = \frac{17}{8(x+2)}$$

$$\frac{1}{(x+2)^{\frac{3}{2}}} - \frac{17}{8(x+2)} = -\frac{1}{4},$$

$$\frac{1}{(x+2)^{\frac{3}{2}}} - \frac{17}{8(x+2)} + \frac{289}{256} = \frac{289}{256} - \frac{1}{4} = \frac{225}{256}$$

$$\frac{1}{x+2} - \frac{17}{16} = \pm \frac{15}{16}$$

$$\frac{1}{x+2} = 2 \text{ or } \frac{1}{8}$$

$$x + 2 = 8 \text{ or } \frac{1}{2}$$

$$x = 6 \text{ or } -\frac{3}{2}.$$

66. Given $\frac{x}{x+4} + \frac{4}{\sqrt{x+4}} = \frac{21}{x}$, to find the values of x .

Then $\frac{x^2}{x+4} + \frac{4x}{\sqrt{x+4}} = 21$

$$\frac{x^2}{x+4} + \frac{4x}{\sqrt{x+4}} + 4 = 25$$

$$\frac{x}{\sqrt{x+4}} + 2 = \pm 5$$

$$\frac{x}{\sqrt{x+4}} = 3 \text{ or } -7$$

$$\frac{x^2}{x+4} = 9 \text{ or } 49.$$

$$\therefore x^2 = 9x + 36$$

$$\text{or } x^2 = 49x + 196$$

$$x^2 - 9x + \frac{81}{4} = \frac{225}{4}$$

$$x^2 - 49x + \frac{2401}{4} = \frac{3185}{4}$$

$$x - \frac{9}{2} = \pm \frac{15}{2}$$

$$x - \frac{49}{2} = \pm \frac{\sqrt{3185}}{2}$$

$$x = 12 \text{ or } -3, \text{ or } \frac{49 \pm \sqrt{3185}}{2}.$$

67. Given $\frac{3x+5}{3x-5} - \frac{3x-5}{3x+5} = \frac{135}{176}$, to find the values of x .

then $\frac{9x^2 + 30x + 25 - 9x^2 + 30x - 25}{9x^2 - 25} = \frac{135}{176},$

$$\text{or } \frac{60x}{9x^2 - 25} = \frac{135}{176};$$

$$\therefore \frac{4x}{9x^2 - 25} = \frac{9}{176},$$

$$81x^2 - 225 = 704x,$$

$$81x^2 - 704x = 225,$$

$$x^2 - \frac{704}{81}x = \frac{225}{81},$$

$$\begin{aligned} x^2 - \frac{704}{81}x + \frac{123904}{6561} &= \frac{123904}{6561} + \frac{225}{81} = \frac{123904 + 18225}{6561} \\ &= \frac{142129}{6561} \end{aligned}$$

$$\therefore x - \frac{352}{81} = \pm \frac{377}{81}$$

$$\text{and } x = 9 \text{ or } -\frac{25}{81}.$$

68. Given $x + \sqrt{x} + 2 = \frac{x^3 + x - 4}{\sqrt{x}}$, to find the values of x .

$$\text{Then } x + \sqrt{x} + 2 = \frac{x^3 - 4}{\sqrt{x}} + \sqrt{x};$$

$$\therefore x + 2 = \frac{x^3 - 4}{\sqrt{x}},$$

$$\text{and } \sqrt{x} = x - 2,$$

$$x - \sqrt{x} = 2,$$

$$x - \sqrt{x} + \frac{1}{4} = \frac{9}{4}$$

$$\sqrt{x} - \frac{1}{2} = \pm \frac{3}{2},$$

$$\sqrt{x} = 2 \text{ or } -1$$

$$\therefore x = 4 \text{ or } 1.$$

69. Given $\frac{x^2}{x^2-4} + \frac{6}{x^2-4} = \frac{351}{25x^2}$, to find the values of x .

$$\text{Then } \frac{25x^4}{x^2-4} + \frac{30 \times 5x^2}{x^2-4} + 225 = 351 + 225 = 576,$$

$$\frac{5x^2}{x^2-4} + 15 = \pm 24,$$

$$\frac{5x^2}{x^2-4} = 9 \text{ or } -39;$$

$$\therefore 5x^2 = 9x^2 - 36$$

$$\text{or } 5x^2 = -39x^2 + 156$$

$$4x^2 = 36$$

$$44x^2 = 156$$

$$x^2 = 9$$

$$x^2 = \frac{156}{44} = \frac{39}{11}$$

$$x = \pm 3$$

$$x = \pm \sqrt{\frac{39}{11}}.$$

70. Given $x + \frac{8}{x} + x = 42 - \frac{8}{x}$, to find the values of x .

$$\text{Here } x + \frac{8}{x} + \left(x + \frac{8}{x}\right) + \frac{1}{4} = 42 + \frac{1}{4} = \frac{169}{4}.$$

$$x + \frac{8}{x} + \frac{1}{2} = \pm \frac{13}{2}$$

$$x + \frac{8}{x} = 6 \text{ or } -7:$$

$$\therefore x^2 + 8 = 6x$$

$$\text{or } x^2 + 8 = -7x$$

$$x^2 - 6x = -8$$

$$x^2 + 7x = -8$$

$$x^2 - 6x + 9 = 1$$

$$x^2 + 7x + \frac{49}{4} = \frac{49}{4} - \frac{32}{4} = \frac{17}{4}$$

$$x - 3 = \pm 1$$

$$\therefore x = 4 \text{ or } 2$$

$$x + \frac{7}{2} = \frac{\pm \sqrt{17}}{2}$$

$$\therefore x = \frac{-7 \pm \sqrt{17}}{2}.$$

71. Given $x + 4 - 2\sqrt{\frac{x+4}{x-4}} = \frac{3}{x-4}$, to find the values of x .

$$(x^2 - 16) - 2\sqrt{x^2 - 16} = 3,$$

$$(x^2 - 16) - 2\sqrt{x^2 - 16} + 1 = 4,$$

$$\sqrt{x^2 - 16} - 1 = \pm 2.$$

$$\sqrt{x^2 - 16} = 3 \text{ or } -1,$$

$$x^2 - 16 = 9 \text{ or } 1,$$

$$x^2 = 25 \text{ or } 17;$$

$$\therefore x = \pm 5 \text{ or } \pm \sqrt{17}$$

72. Given $x^4 \left(1 + \frac{1}{3x}\right)^2 - (3x^2 + x) = 70$, to find the values of x .

$$x^4 \left(1 + \frac{1}{3x}\right)^2 - 3x^2 \left(1 + \frac{1}{3x}\right) + \frac{9}{4} = 70 + \frac{9}{4} = \frac{289}{4}$$

$$x^2 \left(1 + \frac{1}{3x}\right) - \frac{3}{2} = \pm \frac{17}{2},$$

$$x^2 \left(1 + \frac{1}{3x}\right) = 10 \text{ or } -7;$$

$$\therefore x^2 + \frac{x}{3} = 10 \text{ or } -7,$$

$$x^2 + \frac{x}{3} + \frac{1}{36} = \frac{361}{36} \text{ or } -\frac{251}{36},$$

$$x + \frac{1}{6} = \pm \frac{19}{6} \text{ or } \pm \frac{\sqrt{251}}{6},$$

$$x = 3 \text{ or } -\frac{10}{3} \text{ or } \frac{-1 \pm \sqrt{-251}}{6}.$$

73. Given $x^3 - \frac{5x}{2} + 15 = \frac{25x^2}{16} - \frac{64}{x^2}$, to find the values of x .

$$x^2 15 + \frac{64}{x^2} = \frac{25x^2}{16} + \frac{5x}{2}$$

$$\text{then } x^2 + 16 + \frac{64}{x^2} = \frac{25x^2}{16} + \frac{5x}{2} + 1,$$

$$x + \frac{8}{x} = \pm \left(\frac{5x}{4} + 1 \right),$$

$$\text{or } 4x^2 + 32 = -5x^2 - 4x \frac{x}{4} + 1 = \frac{8}{x}$$

$$x^2 + 4x = 32$$

$$9x^2 + 4x = -32$$

$$x^2 + 4x + 4 = 36$$

$$x^2 + \frac{4x}{9} + \frac{4}{81} = \frac{4}{81} - 32 = \frac{-284}{81}$$

$$x + 2 = \pm 6$$

$$x + \frac{2}{9} = \pm \frac{\sqrt{-284}}{9} \text{ or } \frac{\pm 2 \sqrt{-71}}{9}$$

$$x = 4 \text{ or } -8 \text{ or } \frac{-2 \pm 2 \sqrt{-71}}{9}$$

74. Given $\frac{35^{\frac{1}{2}}}{\sqrt{x^2 - 9x}} + \frac{\sqrt{x^2 - 9}}{7x} = \frac{19}{2x}$, to find the values of x .

$$\text{or } \frac{250}{7\sqrt{x^2 - 9}} + \frac{\sqrt{x^2 - 9}}{7} = \frac{19}{2},$$

$$250 + (x^2 - 9) = \frac{133\sqrt{x^2 - 9}}{2},$$

$$(x^2 - 9) - \frac{133\sqrt{x^2 - 9}}{2} + \frac{133^2}{4} = \frac{17689}{16} - 250 = \frac{17689 - 4000}{16} = \frac{13689}{16}$$

$$\sqrt{x^2 - 9} - \frac{133}{4} = \frac{\pm 117}{4},$$

$$\sqrt{x^2 - 9} = \frac{16}{4} \text{ or } \frac{250}{4}$$

$$= 4 \text{ or } \frac{125}{2}$$

$$x^2 - 9 = 16 \text{ or } \frac{15625}{4}$$

$$x^2 = 25 \text{ or } \frac{15661}{4}$$

$$x = \pm 5 \text{ or } \pm \frac{\sqrt{15661}}{2}$$

75. Given $3.\overline{x-1}^2 - x + 2x = 341 + 2.\overline{x-1}^2$, to find the values of x .

$$\text{Then } 3.\overline{x-1}^2 - x - 2.\overline{x-1}^2 - x = 341,$$

$$\overline{x-1}^2 - x = \frac{2}{3}.\overline{x-1}^2 - x + \frac{1}{9} = \frac{341}{3} + \frac{1}{9} = \frac{1024}{9}$$

$$(\overline{x-1}^2 - x) - \frac{1}{9} = \pm \frac{32}{3}$$

$$\overline{x-1}^2 - x = 11 \text{ or } -\frac{31}{3}$$

$$x^2 - 2x + 1 - x = 11 \text{ or } -\frac{31}{3}$$

$$x^2 - 3x = 10 \text{ or } -\frac{34}{3}$$

$$x^2 - 3x + \frac{9}{4} = \frac{49}{4} \text{ or } \frac{-109}{12}$$

$$x - \frac{3}{2} = \pm \frac{7}{2} \text{ or } \frac{\pm \sqrt{-109}}{2\sqrt{3}}$$

$$\therefore x = 5 \text{ or } -2 \text{ or } \frac{3}{2} \pm \frac{\sqrt{-109}}{2\sqrt{3}} = \frac{3\sqrt{3} \pm \sqrt{-109}}{2\sqrt{3}}$$

76. Given $x^4 + \frac{13}{3}x^2 - 39x = 81$, to find the values of x .

$$\text{Here } x^4 + \frac{13}{3}x^2 = 39x + 81,$$

$$\therefore x^4 + \frac{13x^3}{3} + \frac{169x^2}{36} = \frac{169x^2}{36} + 39x + 81,$$

$$x^2 + \frac{13x}{6} = \pm \left(\frac{13x}{6} + 9 \right),$$

and first, $x^2 = 9$,

$$\therefore x = \pm 3,$$

secondly, $x^2 + \frac{26x}{6} = -9$,

$$x^2 + \frac{26x}{6} + \frac{169}{36} = \frac{169}{36} - 9 = \frac{169 - 324}{36} = \frac{-155}{36}$$

$$x + \frac{13}{6} = \frac{\pm \sqrt{-155}}{6}$$

$$x = \frac{-13 \pm \sqrt{-155}}{6}.$$

77. Given $4x + \frac{x}{2} = 4x^3 + 33$, to find the values of x .

$$\text{Here } 4x^4 - 4x^3 = 33 - \frac{x}{2}$$

$$4x^4 - 4x^3 + x^2 = 33 + x^2 - \frac{x}{2} = 33 + \frac{2x^2 - x}{2}$$

$$\overline{2x^2 - x}^2 - \frac{2x^2 - x}{2} + \frac{1}{16} = 33 + \frac{1}{16} = \frac{529}{16}$$

$$2x^2 - x - \frac{1}{4} = \pm \frac{23}{4}$$

$$2x^2 - x = 6 \text{ or } -\frac{11}{2}$$

$$x^2 - \frac{x}{2} = 3 \text{ or } -\frac{11}{4}$$

$$x^2 - \frac{x}{2} + \frac{1}{16} = \frac{49}{16} \text{ or } -\frac{43}{16}$$

$$x - \frac{1}{4} = \pm \frac{7}{4} \text{ or } \frac{\pm \sqrt{-43}}{4}$$

$$\text{and } x = 2 \text{ or } -\frac{3}{2} \text{ or } \frac{1 \pm \sqrt{-43}}{4}.$$

78. Given $\overline{x-2}^2 - 6x^{\frac{1}{2}}(x-2) = 24 - 14x + 15x^{\frac{1}{2}}$
to find the values of x .

$$\overline{x-2}^2 - 6x^{\frac{1}{2}}(x-2) + 9x = 24 - 5x + 15x^{\frac{1}{2}},$$

$$\therefore x - 2 - 3x^{\frac{1}{2}} = \pm \sqrt{24 - 5(x - 3x^{\frac{1}{2}})}$$

$$\text{or } (x - 3x^{\frac{1}{2}}) - 2 = \pm \sqrt{24 - 5(x - 3x^{\frac{1}{2}})},$$

$$\therefore (x - 3x^{\frac{1}{2}})^2 - 4(x - 3x^{\frac{1}{2}}) + 4 = 24 - 5(x - 3x^{\frac{1}{2}})$$

$$(x - 3x^{\frac{1}{2}})^2 + (x - 3x^{\frac{1}{2}}) = 20,$$

$$(x - 3x^{\frac{1}{2}})^2 + (x - 3x^{\frac{1}{2}}) + \frac{1}{4} = \frac{81}{4},$$

$$\therefore (x - 3x^{\frac{1}{2}}) + \frac{1}{2} = \pm \frac{9}{2}$$

$$\text{and } x - 3x^{\frac{1}{2}} = 4 \text{ or } -5,$$

$$\text{Again, } x - 3x^{\frac{1}{2}} + \frac{9}{4} = \frac{25}{4} \text{ or } -\frac{11}{4},$$

$$\therefore x^{\frac{1}{2}} - \frac{3}{2} = \pm \frac{5}{2} \text{ or } \pm \frac{\sqrt{-11}}{2};$$

$$x^{\frac{1}{2}} = 4 \text{ or } -1 \text{ or } \frac{3 \pm \sqrt{-11}}{2}$$

$$\text{and } x = 16 \text{ or } 1 \text{ or } \frac{9 \pm 6\sqrt{-11} - 11}{4}$$

$$\text{or } x = 16 \text{ or } 1 \text{ or } \frac{\pm 3\sqrt{-11} - 1}{2}$$

79. Given $\overline{4x+1}^2 + 4x^{\frac{1}{2}}(4x+1) = 1912 - (10x + 3x^{\frac{1}{2}})$,
to find the values of x .

$$\overline{4x+1}^2 + 4x^{\frac{1}{2}}(4x+1) + 4x = 1912 - (6x + 3x^{\frac{1}{2}}),$$

$$\therefore 4x + 1 + 2x^{\frac{1}{2}} = \sqrt{1912 - 3(2x + x^{\frac{1}{2}})}$$

$$\begin{aligned} & \text{or } 2.(2x+x^{\frac{1}{2}}) + 1 = \sqrt{1912-3.(2x+x^{\frac{1}{2}})} \\ \therefore 4(2x+x^{\frac{1}{2}})^2 + 4(2x+x^{\frac{1}{2}}) + 1 &= 1912-3.(2x+x^{\frac{1}{2}}), \\ \text{and } 2x+x^{\frac{1}{2}} + \frac{7}{4}(2x+x^{\frac{1}{2}}) &= \frac{1911}{4}, \end{aligned}$$

$$2x+x^{\frac{1}{2}} + \frac{7}{4}(2x+x^{\frac{1}{2}}) + \frac{49}{64} = \frac{1911}{4} + \frac{49}{64} = \frac{30625}{64},$$

$$\therefore 2x+x^{\frac{1}{2}} + \frac{7}{8} = \pm \frac{175}{8}$$

$$\text{and } 2x+x^{\frac{1}{2}} = 21 \text{ or } -\frac{182}{8},$$

$$\text{or } x + \frac{x^{\frac{1}{2}}}{2} = \frac{21}{2} \text{ or } -\frac{91}{8}$$

$$\text{Again } x + \frac{x^{\frac{1}{2}}}{2} + \frac{1}{16} = \frac{168+1}{16} \text{ or } \frac{1-182}{16},$$

$$= \frac{169}{16} \text{ or } \frac{-181}{16},$$

$$\therefore x^{\frac{1}{2}} + \frac{1}{4} = \pm \frac{13}{4} \text{ or } \pm \frac{\sqrt{-181}}{4},$$

$$x^{\frac{1}{2}} = 3 \text{ or } -\frac{7}{2} \text{ or } \frac{-1 \pm \sqrt{-181}}{4}$$

$$\text{and } x = 9 \text{ or } \frac{49}{4} \text{ or } \frac{-90 \mp \sqrt{-181}}{8}.$$

80. Given $8x^2 - 13 = \frac{3x}{2} + \sqrt{6x^2 + 52x^2}$, to find the values of x .

Multiplying by 4 we have,

$$32x^2 - 52 = 6x + 4x\sqrt{6x+52},$$

$$\text{or } 6x + 52 + 4x\sqrt{6x+52} + 4x^2 = 32x^2 + 4x^2 = 36x^2,$$

$$\sqrt{6x+52} + 2x = \pm 6x,$$

$$\sqrt{6x+52} = 4x \text{ or } -8x,$$

$$6x+52 = 16x^2 \text{ or } 64x^2;$$

$$\begin{aligned}
 \therefore 16x^2 - 6x &= 52 & \text{or } 64x^2 - 6x &= 52, \\
 x^2 - \frac{3x}{8} &= \frac{13}{4} & x^2 - \frac{3x}{32} &= \frac{13}{16} \\
 x^2 - \frac{3x}{8} + \frac{9}{256} &= \frac{841}{256} & x^2 - \frac{3x}{32} + \frac{9}{4096} &= \frac{3337}{4096} \\
 x - \frac{3}{16} &= \pm \frac{29}{16} & x - \frac{3}{64} &= \pm \frac{\sqrt{3337}}{64} \\
 x &= 2 \text{ or } -\frac{13}{8} & \text{or } \frac{3 \pm \sqrt{3337}}{64}.
 \end{aligned}$$

81. Given $4x^2 + 21x + 8x^{\frac{1}{2}}\sqrt{7x^2 - 5} = 207 - \frac{4x^2}{3}$,
to find the values of x .

$$\begin{aligned}
 \text{Here } \frac{16x^2}{3} + 21x + 8x\sqrt{7x^2 - 5} &= 207, \\
 \frac{16x^2}{9} + 7x + \frac{8x}{3}\sqrt{7x^2 - 5} &= 69, \\
 7x - 5 + \frac{8x}{3}\sqrt{7x^2 - 5} + \frac{16x^2}{9} &= 64 \\
 \sqrt{7x^2 - 5} + \frac{4x}{3} &= \pm 8, \\
 3\sqrt{7x^2 - 5} &= \pm 24 - 4x, \\
 63x - 45 &= 576 \mp 192x + 16x^2; \\
 \therefore 16x^2 - 255x &= -621, & \text{or } 16x^2 + 129x &= -621 \\
 x^2 - \frac{255}{16} + \frac{65025}{1024} &= \frac{65025}{1024} - \frac{621}{16} & x^2 + \frac{129}{16}x + \frac{129^2}{32} &= \frac{16641}{1024} - \frac{621}{16} \\
 & & &= \frac{16641 - 39744}{1024}, \\
 & & &= \frac{25281}{1024} & &= \frac{-23103}{1024};
 \end{aligned}$$

$$x - \frac{255}{32} = \pm \frac{159}{32} \quad x + \frac{129}{32} = \pm \frac{\sqrt{-23103}}{32}$$

$$x = \frac{96}{32} \text{ or } \frac{414}{32} = \pm \frac{3\sqrt{-2567}}{32},$$

$$x = 3 \text{ or } \frac{207}{16} \quad \text{or } \frac{-129 \pm 3\sqrt{-2567}}{32}.$$

82. Given $\frac{2x + \sqrt{x}}{2x - \sqrt{x}} = 3 \frac{7}{15} - 3$, $\frac{2x - \sqrt{x}}{2x + \sqrt{x}}$, to find the values of x .

$$\frac{2\sqrt{x+1}}{2\sqrt{x-1}} + 3 = \frac{2\sqrt{x-1}}{2\sqrt{x+1}} = \frac{52}{15}$$

$$\frac{4x + 4\sqrt{x+1} + 12x - 12\sqrt{x+1} + 3}{4x-1} = \frac{52}{15}$$

$$\frac{16x - 8\sqrt{x+1} + 4}{4x-1} = \frac{52}{15}$$

$$\frac{4x - 2\sqrt{x+1}}{4x-1} = \frac{13}{15}$$

$$60x - 30\sqrt{x+1} + 15 = 52x - 13,$$

$$8x - 30\sqrt{x+1} = -28,$$

$$x - \frac{15}{4}\sqrt{x+1} = -\frac{14}{4},$$

$$x - \frac{15}{4}\sqrt{x+1} + \frac{225}{64} = \frac{225}{64} - \frac{14}{4} = \frac{1}{64}.$$

$$\sqrt{x+1} - \frac{15}{8} = \pm \frac{1}{8},$$

$$\sqrt{x+1} = 2 \text{ or } \frac{7}{4},$$

$$x = 4 \text{ or } \frac{49}{16}.$$

$$\sqrt{y} - \frac{1}{4} = \pm \frac{7}{4} \text{ or } \frac{\pm \sqrt{241}}{4}$$

$$\sqrt{y} = 2 \text{ or } -\frac{3}{2} \text{ or } \frac{1 \pm \sqrt{241}}{4}$$

$$y = 4 \text{ or } \frac{9}{4} \text{ or } \frac{1 \pm 2\sqrt{241} + 241}{16}$$

$$= 4 \text{ or } \frac{9}{4} \text{ or } \frac{242 \pm 2\sqrt{241}}{16}$$

$$= 4 \text{ or } \frac{9}{4} \text{ or } \frac{121 \pm \sqrt{241}}{8}$$

$$\text{and } x = 5 \text{ or } \frac{17}{2} \text{ or } \frac{-69 \mp \sqrt{241}}{4}$$

11. Given $\frac{x^4}{y^2} + \frac{2x^2}{y} = 9\frac{39}{49}$ } to find the values of x
 and $x + y^2 = 65$ } and y .

$$\frac{x^4}{y^2} + \frac{2x^2}{y} = \frac{480}{49},$$

$$\frac{x^4}{y^2} + \frac{2x^2}{y} + 1 = \frac{480}{49} + 1 = \frac{529}{49},$$

$$\frac{x^2}{y} + 1 = \pm \frac{23}{7},$$

$$\frac{x^2}{y} = \pm \frac{23}{7} - 1 = \frac{16}{7} \text{ or } -\frac{30}{7},$$

$$x^2 = \frac{16y}{7} \text{ or } -\frac{30y}{7},$$

$$y^2 + \frac{16y}{7} + \frac{64}{49} = 65 + \frac{64}{49} = \frac{3249}{49},$$

$$y + \frac{8}{7} = \pm \frac{57}{7},$$

$$y = 7 \text{ or } -\frac{65}{7},$$

9. Given $\overline{x+y}^2 - 3y = 28 + 3x$ } to find the values of
and $2xy + 3x = 35$ } x and y .

$$\overline{x+y}^2 - 3 \cdot \overline{x+y} + \frac{9}{4} = 28 + \frac{9}{4} = \frac{121}{4}$$

$$x + y - \frac{3}{2} = \pm \frac{11}{2}$$

$$x + y = 7 \text{ or } -4$$

$$y = 7 - x \text{ or } -x - 4;$$

$$\therefore 14x - 2x^2 + 3x = 35 \quad \text{or} \quad -2x^2 - 8x + 3x = 35$$

$$2x^2 - 17x = -35 \quad 2x^2 + 5x = -35$$

$$x^2 - \frac{17x}{2} + \frac{289}{16} = \frac{9}{16} \quad x^2 + \frac{5x}{2} + \frac{25}{16} = -\frac{255}{16}$$

$$x - \frac{17}{4} = \pm \frac{3}{4} \quad x + \frac{5}{4} = \frac{\pm \sqrt{-255}}{4}$$

$$x = 5 \text{ or } \frac{7}{2} \quad x = \frac{-5 \pm \sqrt{-255}}{4}$$

$$\therefore x = 5 \text{ or } \frac{7}{2} \text{ or } \frac{-5 \pm \sqrt{-255}}{4}; \text{ and } y = 2 \text{ or } \frac{7}{2} \text{ or } \frac{-11 \mp \sqrt{-255}}{4}.$$

10. Given $x^2 + 10x + y = 119 - 2\sqrt{y} \cdot (x + 5)$ }
and $x + 2y = 13$ }

to find the values of x and y .

$$x^2 + 2x\sqrt{y} + y + 10(x + \sqrt{y}) + 25 = 144,$$

$$x + \sqrt{y} + 5 = \pm 12,$$

$$\therefore x + \sqrt{y} = 7 \text{ or } -17$$

$$\text{but } x + 2y = 13$$

$$\therefore 2y - \sqrt{y} = 6 \text{ or } 30$$

$$y - \frac{\sqrt{y}}{2} = 3 \text{ or } 15$$

$$y - \frac{\sqrt{y}}{2} + \frac{1}{16} = \frac{49}{16} \text{ or } \frac{241}{16}$$

$$\sqrt{y} - \frac{1}{4} = \pm \frac{7}{4} \text{ or } \frac{\pm \sqrt{241}}{4}$$

$$\sqrt{y} = 2 \text{ or } -\frac{3}{2} \text{ or } \frac{1 \pm \sqrt{241}}{4}$$

$$y = 4 \text{ or } \frac{9}{4} \text{ or } \frac{1 \pm 2\sqrt{241} + 241}{16}$$

$$= 4 \text{ or } \frac{9}{4} \text{ or } \frac{242 \pm 2\sqrt{241}}{16}$$

$$= 4 \text{ or } \frac{9}{4} \text{ or } \frac{121 \pm \sqrt{241}}{8}$$

$$\text{and } x = 5 \text{ or } \frac{17}{2} \text{ or } \frac{-69 \mp \sqrt{241}}{4}$$

11. Given $\frac{x^4}{y^2} + \frac{2x^2}{y} = 9\frac{39}{49}$ } to find the values of x
 and $x + y^2 = 65$ } and y .

$$\frac{x^4}{y^2} + \frac{2x^2}{y} = \frac{480}{49},$$

$$\frac{x^4}{y^2} + \frac{2x^2}{y} + 1 = \frac{480}{49} + 1 = \frac{529}{49},$$

$$\frac{x^2}{y} + 1 = \pm \frac{23}{7},$$

$$\frac{x^2}{y} = \pm \frac{23}{7} - 1 = \frac{16}{7} \text{ or } -\frac{30}{7},$$

$$x^2 = \frac{16y}{7} \text{ or } -\frac{30y}{7},$$

$$y^2 + \frac{16y}{7} + \frac{64}{49} = 65 + \frac{64}{49} = \frac{3249}{49},$$

$$y + \frac{8}{7} = \pm \frac{57}{7},$$

$$y = 7 \text{ or } -\frac{65}{7},$$

$$\text{Again, } y^2 - \frac{30y}{7} = 65,$$

$$y^2 - \frac{30y}{7} + \frac{225}{49} = 65 + \frac{225}{49},$$

$$= \frac{3410}{49},$$

$$y - \frac{15}{7} = \frac{\pm \sqrt{3410}}{7},$$

$$y = \frac{15 \pm \sqrt{3410}}{7},$$

$$\therefore x = \pm 4 \text{ or } \frac{\pm 4 \sqrt{-65}}{7} \text{ or } \frac{\pm \sqrt{-450 \mp 30 \sqrt{3410}}}{\sqrt{7}}.$$

12. Given $x + y + \sqrt{x + y} = 6$ } to find the values
and $x^2 + y^2 = 10$ } of x and y .

$$x + y + \sqrt{x + y} + \frac{1}{4} = \frac{25}{4},$$

$$\sqrt{x + y} + \frac{1}{2} = \pm \frac{5}{2},$$

$$\sqrt{x + y} = 2 \text{ or } -3,$$

$$x + y = 4 \text{ or } 9,$$

$$\therefore x^2 + 2xy + y^2 = 16 \text{ or } 81,$$

$$\text{but } x^2 + y^2 = 10,$$

$$\therefore 2xy = 6 \text{ or } 71,$$

$$x^2 - 2xy + y^2 = 4 \text{ or } -61.$$

$$x - y = \pm 2 \text{ or } \pm \sqrt{-61},$$

$$x + y = 4 \text{ or } 9.$$

$$2x = 6 \text{ or } 2 \text{ or } 9 \pm \sqrt{-61},$$

$$x = 3 \text{ or } 1 \text{ or } \frac{9 \pm \sqrt{-61}}{2},$$

$$y = 1 \text{ or } 3 \text{ or } \frac{9 \mp \sqrt{-61}}{2},$$

13. Given $x^2 + 4\sqrt{x^2 + 3y + 5} = 55 - 3y$ } to find the
and $6x - 7y = 16$ } values of x and y .

$$x^2 + 3y + 5 + 4\sqrt{x^2 + 3y + 5} + 4 = 64,$$

$$\therefore \sqrt{x^2 + 3y + 5} + 2 = \pm 8,$$

$$\text{and } \sqrt{x^2 + 3y + 5} = 6 \text{ or } -10,$$

$$x^2 + 3y + 5 = 36 \text{ or } 100,$$

$$x^2 + 3y = 31 \text{ or } 95,$$

$$7x^2 + 21y = 217 \text{ or } 665,$$

$$18x - 21y = 48,$$

$$\therefore 7x^2 + 18x = 265 \text{ or } 713,$$

$$x^2 + \frac{18x}{7} + \frac{81}{49} = \frac{265}{7} + \frac{81}{49} \text{ or } \frac{713}{7} + \frac{81}{49},$$

$$= \frac{1936}{49} \text{ or } \frac{5072}{49},$$

$$x + \frac{9}{7} = \pm \frac{44}{7} \text{ or } \frac{\pm \sqrt{5072}}{7},$$

$$x = 5 \text{ or } \frac{-53}{7} \text{ or } \frac{-9 \pm 4\sqrt{317}}{7},$$

$$\therefore y = 2 \text{ or } \frac{430}{49} \text{ or } \frac{-166 \mp 24\sqrt{317}}{49}.$$

14. Given $x^2 + 3x + y = 73 - 2xy$ } to find the values
and $y^2 + 3y + x = 44$ } of x and y .

$$x^2 + 2xy + y^2 + 4x + 4y = 117,$$

$$x + y^2 + 4x + y + 4 = 121,$$

$$x + y + 2 = \pm 11,$$

$$x + y = 9 \text{ or } -13,$$

$$\text{but } y^2 + x + 3y = 44,$$

$$\therefore y^2 + 2y = 35 \text{ or } 57,$$

$$y^2 + 2y + 1 = 36 \text{ or } 58,$$

$$\begin{aligned} & \text{or } 2.(2x+x^{\frac{1}{2}}) + 1 = \sqrt{1912-3.(2x+x^{\frac{1}{2}})} \\ \therefore 4(2x+x^{\frac{1}{2}})^2 + 4(2x+x^{\frac{1}{2}}) + 1 &= 1912-3.(2x+x^{\frac{1}{2}}), \\ \text{and } 2x+x^{\frac{1}{2}} + \frac{7}{4}(2x+x^{\frac{1}{2}}) &= \frac{1911}{4}, \end{aligned}$$

$$2x+x^{\frac{1}{2}} + \frac{7}{4}(2x+x^{\frac{1}{2}}) + \frac{49}{64} = \frac{1911}{4} + \frac{49}{64} = \frac{30625}{64},$$

$$\therefore 2x+x^{\frac{1}{2}} + \frac{7}{8} = \pm \frac{175}{8}$$

$$\text{and } 2x+x^{\frac{1}{2}} = 21 \text{ or } -\frac{182}{8},$$

$$\text{or } x + \frac{x^{\frac{1}{2}}}{2} = \frac{21}{2} \text{ or } -\frac{91}{8}$$

$$\begin{aligned} \text{Again } x + \frac{x^{\frac{1}{2}}}{2} + \frac{1}{16} &= \frac{168+1}{16} \text{ or } \frac{1-182}{16}, \\ &= \frac{169}{16} \text{ or } \frac{-181}{16}, \end{aligned}$$

$$\therefore x^{\frac{1}{2}} + \frac{1}{4} = \pm \frac{13}{4} \text{ or } \pm \frac{\sqrt{-181}}{4},$$

$$x^{\frac{1}{2}} = 3 \text{ or } -\frac{7}{2} \text{ or } \frac{-1 \pm \sqrt{-181}}{4}$$

$$\text{and } x = 9 \text{ or } \frac{49}{4} \text{ or } \frac{-90 \mp \sqrt{-181}}{8}.$$

80. Given $8x^2 - 13 = \frac{3x}{2} + \sqrt{6x^3 + 52x^2}$, to find the values of x .

Multiplying by 4 we have,

$$\begin{aligned} 32x^2 - 52 &= 6x + 4x\sqrt{6x+52}, \\ \text{or } 6x + 52 + 4x\sqrt{6x+52} + 4x^2 &= 32x^2 + 4x^2 = 36x^2, \\ \sqrt{6x+52} + 2x &= \pm 6x, \\ \sqrt{6x+52} &= 4x \text{ or } -8x, \\ 6x+52 &= 16x^2 \text{ or } 64x^2; \end{aligned}$$

$$\therefore 16x^2 - 6x = 52$$

$$\text{or } 64x^2 - 6x = 52,$$

$$x^2 - \frac{3x}{8} = \frac{13}{4}$$

$$x^2 - \frac{3x}{32} = \frac{13}{16}$$

$$x^2 - \frac{3x}{8} + \frac{9}{256} = \frac{841}{256}$$

$$x^2 - \frac{3x}{32} + \frac{9}{4096} = \frac{3337}{4096}$$

$$x - \frac{3}{16} = \pm \frac{29}{16}$$

$$x - \frac{3}{64} = \pm \frac{\sqrt{3337}}{64}$$

$$x = 2 \text{ or } -\frac{13}{8} \text{ or } \frac{3 \pm \sqrt{3337}}{64}.$$

$$81. \text{ Given } 4x^2 + 21x + 8x^{\frac{1}{2}}\sqrt{7x^2 - 5} = 207 - \frac{4x^2}{3},$$

to find the values of x .

$$\text{Here } \frac{16x^2}{3} + 21x + 8x\sqrt{7x-5} = 207,$$

$$\frac{16x^2}{9} + 7x + \frac{8x}{3}\sqrt{7x-5} = 69,$$

$$7x-5 + \frac{8x}{3}\sqrt{7x-5} + \frac{16x^2}{9} = 64$$

$$\sqrt{7x-5} + \frac{4x}{3} = \pm 8,$$

$$3\sqrt{7x-5} = \pm 24 - 4x,$$

$$63x - 45 = 576 \mp 192x + 16x^2;$$

$$\therefore 16x^2 - 255x = -621, \quad \text{or } 16x^2 + 129x = -621$$

$$x^2 - \frac{255}{16} + \frac{65025}{1024} = \frac{65025}{1024} - \frac{621}{16}$$

$$x^2 + \frac{129}{16}x + \frac{129}{32} \Bigg|^2$$

$$= \frac{16641}{1024} - \frac{621}{16}$$

$$= -\frac{65025-39744}{1024}$$

$$= \frac{16641-39744}{1024},$$

$$= \frac{25281}{1024}$$

$$= \frac{-23103}{1024};$$

17. Given $x^4 + y^4 = 97$ } to find the values of x and y .
 and $x + y = 5$ }

$$\begin{array}{r} x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4 = 625 \\ x^4 \qquad \qquad \qquad + y^4 = 97, \end{array}$$

$$\therefore 4x^3y + 6x^2y^2 + 4xy^3 = 528$$

$$\text{or } 2x^3y + 3x^2y^2 + 2xy^3 = 264$$

$$\therefore 2x^3 + 3xy + 2y^3 = \frac{264}{xy}$$

$$\text{but } 2x^3 + 4xy + 2y^3 = 50$$

$$xy = 50 - \frac{264}{xy},$$

$$x^3y^3 - 50xy = -264,$$

$$x^3y^3 - 50xy + 625 = 625 - 264 = 361,$$

$$xy - 25 = \pm 19,$$

$$xy = 44 \text{ or } 6,$$

$$x^3 + 2xy + y^3 = 25.$$

$$4xy = 24 \text{ or } 176,$$

$$x^3 - 2xy + y^3 = 1 \text{ or } -151,$$

$$\therefore x - y = \pm 1 \text{ or } \pm \sqrt{-151},$$

$$\text{but } x + y = 5,$$

$$\text{consequently } x = 3 \text{ or } 2 \text{ or } \frac{5 \pm \sqrt{-151}}{2},$$

$$\text{and } y = 2 \text{ or } 3 \text{ or } \frac{5 \mp \sqrt{-151}}{2}.$$

18. Given $\sqrt{\frac{3x-2y}{2x}} + \sqrt{\frac{2x}{3x-2y}} = 2$ } to find the
 and $x^2 - 18 = x(4y - 9)$ } values of
 x and y .

Multiplying both sides of the first equation by $\sqrt{\frac{3x-2y}{2x}}$,

$$\text{we have } \frac{3x-2y}{2x} + 1 = 2 \cdot \sqrt{\frac{3x-2y}{2x}}$$

$$\text{or } \frac{3x-2y}{2x} - 2 \cdot \sqrt{\frac{3x-2y}{2x}} + 1 = 0,$$

$$\sqrt{\frac{3x-2y}{2x}} - 1 = 0,$$

$$\frac{3x-2y}{2x} = 1,$$

$$\therefore 3x - 2y = 2x,$$

$$x = 2y,$$

$$\text{and } y = \frac{x}{2};$$

$$\text{Again, } x^2 - 18 = x(2x - 9) = 2x^2 - 9x,$$

$$x^2 - 9x = -18,$$

$$x^2 - 9x + \frac{81}{4} = \frac{81-72}{4} = \frac{9}{4},$$

$$x - \frac{9}{2} = \pm \frac{3}{2},$$

$$x = 6 \text{ or } 3,$$

$$y = \frac{x}{2} = 3 \text{ or } \frac{3}{2}$$

$$19. \text{ Given } \left. \begin{array}{l} x+4\sqrt{x}+4y=21+8\sqrt{y}+4\sqrt{xy} \\ \text{and } \sqrt{x}+\sqrt{y}=6 \end{array} \right\} \text{ to find}$$

the values of x and y .

$$x - 4\sqrt{xy} + 4y + 4\sqrt{x} - 8\sqrt{y} = 21,$$

$$\text{or } (\sqrt{x} - 2\sqrt{y})^2 + 4(\sqrt{x} - 2\sqrt{y}) + 4 = 25,$$

$$\sqrt{x} - 2\sqrt{y} + 2 = \pm 5,$$

$$\therefore \sqrt{x} - 2\sqrt{y} = 3 \text{ or } -7.$$

$$\text{but } \sqrt{x} + \sqrt{y} = 6,$$

$$3\sqrt{y} = 3 \text{ or } 13,$$

$$\sqrt{y} = 1 \text{ or } \frac{13}{3},$$

$$y = 1 \text{ or } \frac{169}{9},$$

$$\sqrt{x} = 6 - \sqrt{y} = 5 \text{ or } \frac{5}{3},$$

$$x = 25 \text{ or } \frac{25}{9}.$$

20. Given $x + y = 5$ } to find the values of
and $(x^3 + y^3) \times (x^3 + y^3) = 455$ } x and y .

$$x^5 + 5x^4y + 10x^3y^2 + 10x^2y^3 + 5xy^4 + y^5 = 3125$$

$$x^5 + x^3y^2 + x^2y^3 + y^5 = 455$$

$$5x^4y + 9x^3y^2 + 9x^2y^3 + 5xy^4 = 2670$$

$$5x^3 + 9x^2y + 9xy^2 + 5y^3 = \frac{2670}{xy}$$

$$\text{but } 5x^3 + 15x^2y + 15xy^2 + 5y^3 = 625$$

$$\therefore 6x^2y + 6xy^2 = 625 - \frac{2670}{xy}$$

$$\text{or } 6xy(x + y) = 625 - \frac{2670}{xy},$$

$$\text{and } 6xy = 125 - \frac{534}{xy};$$

$$\therefore 6x^2y^2 - 125xy = -534,$$

$$x^2y^2 - \frac{125}{6}xy + \frac{15625}{144} = \frac{15625}{144} - \frac{534}{6}$$

$$= \frac{15625 - 12916}{144},$$

$$= \frac{2809}{144},$$

$$xy - \frac{125}{12} = \pm \frac{53}{12}$$

$$xy = \frac{178}{12} \text{ or } \frac{72}{12}$$

$$= \frac{89}{6} \text{ or } 6,$$

$$\begin{aligned}x^2 + 2xy + y^2 &= 25, \\4xy &= 24 \text{ or } \frac{178}{3}.\end{aligned}$$

$$x^2 - 2xy + y^2 = 1 \text{ or } -\frac{103}{3}$$

$$\therefore x - y = \pm 1 \text{ or } \pm \sqrt{-\frac{103}{3}}$$

$$\text{but } x + y = 5.$$

$$x = 3 \text{ or } 2 \text{ or } \frac{5}{2} \pm \frac{1}{2} \sqrt{-\frac{103}{3}}$$

$$x = 2 \text{ or } 3 \text{ or } \frac{5}{2} \mp \frac{1}{2} \sqrt{-\frac{103}{3}}.$$

$$21. \text{ Given } \left. \begin{aligned}x + y - \sqrt{\frac{x+y}{x-y}} &= \frac{6}{x-y} \\ \text{and } x^2 + y^2 &= 41\end{aligned} \right\} \begin{array}{l} \text{to find the va-} \\ \text{lues of } x \\ \text{and } y. \end{array}$$

$$\text{Here } x^2 - y^2 - \sqrt{x^2 - y^2} = 6,$$

$$x^2 - y^2 - \sqrt{x^2 - y^2} + \frac{1}{4} = \frac{25}{4}$$

$$\sqrt{x^2 - y^2} - \frac{1}{2} = \pm \frac{5}{2}$$

$$\sqrt{x^2 - y^2} = 3 \text{ or } -2,$$

$$\therefore x^2 - y^2 = 9 \text{ or } 4$$

$$\text{but } x^2 + y^2 = 41.$$

$$x^2 = 25 \text{ or } \frac{45}{2},$$

$$x = \pm 5 \pm 3 \sqrt{\frac{5}{2}},$$

$$y^2 = 16 \text{ or } \frac{37}{2},$$

$$y = \pm 4 \text{ or } \pm \sqrt{\frac{37}{2}}.$$

$$22. \text{ Given } \left. \begin{aligned} \frac{x^4}{y^3} + \frac{y^4}{x^3} &= 136 \frac{1}{9} - 2xy \\ \text{and } x + 4 &= 14 - y \end{aligned} \right\} \text{ to find the}$$

values of x and y .

$$\text{Here } \frac{x^4}{y^3} + 2xy + \frac{y^4}{x^3} = \frac{1225}{9}$$

$$\therefore \frac{x^3}{y} + \frac{y^3}{x} = \pm \frac{35}{3}$$

$$\text{or } x^3 + y^3 = \pm \frac{35xy}{3},$$

$$\text{but } x + y = 10;$$

$$\therefore x^3 + 3x^2y + 3xy^2 + y^3 = 1000,$$

$$\text{and } x^3 + y^3 = \pm \frac{35xy}{3},$$

$$\therefore 3xy \cdot \overline{x+y} = 1000 \mp \frac{35xy}{3}$$

$$30xy = 1000 \mp \frac{35xy}{3},$$

$$90xy \pm 35xy = 3000,$$

$$125xy \text{ or } 55xy = 3000$$

$$xy = \frac{3000}{125} \text{ or } \frac{3000}{55}$$

$$= 24 \text{ or } \frac{600}{11}.$$

$$\text{Now } x^3 + 2xy + y^3 = 100$$

$$\text{and } 4xy = 96 \text{ or } \frac{2400}{11},$$

$$\therefore x^3 - 2xy + y^3 = 4 \text{ or } -\frac{1300}{11}$$

$$x - y = \pm 2 \text{ or } \pm 10. \quad \sqrt{-\frac{13}{11}}$$

$$x + y = 10.$$

$$x = 6 \text{ or } 4 \text{ or } 5 \pm 5. \sqrt{-\frac{13}{11}}$$

$$y = 4 \text{ or } 6 \text{ or } 5 \mp 5. \sqrt{-\frac{13}{11}}.$$

$$\left. \begin{array}{l} 23. \text{ Given } \frac{x+y}{x-y} - \frac{x-y}{x+y} = 4 \frac{4}{5} \\ \text{and } \sqrt{\frac{x+y}{x^2}} + \frac{1}{x} = \frac{4}{9\sqrt{x-y}} \end{array} \right\} \text{to find the values of } x \text{ and } y.$$

From the first equation,

$$\frac{x^2 + 2xy + y^2 - x^2 + 2xy - y^2}{x^2 - y^2} = \frac{24}{5};$$

$$\text{or } \frac{4xy}{x^2 - y^2} = \frac{24}{5}.$$

$$\text{Again, } \frac{x-y}{x^2} + \frac{\sqrt{x-y}}{x} = \frac{4}{9}.$$

$$\frac{x-y}{x^2} + \frac{\sqrt{x-y}}{x} + \frac{1}{4} = \frac{1}{4} + \frac{4}{9} = \frac{25}{36};$$

$$\therefore \frac{\sqrt{x-y}}{x} + \frac{1}{2} = \pm \frac{5}{6},$$

$$\text{and } \frac{\sqrt{x-y}}{x} = \frac{1}{3} \text{ or } -\frac{4}{3};$$

$$\therefore \frac{x-y}{x^2} = \frac{1}{9} \text{ or } \frac{16}{9}.$$

Multiplying this equation by $\frac{4xy}{x^2 - y^2} = \frac{24}{5}$

$$\frac{y}{x \cdot x+y} = \frac{2}{15} \text{ or } \frac{32}{15};$$

$$\text{First, } \frac{9-x}{18x-x^2} = \frac{2}{15},$$

$$\text{or } 135 - 15x = 36x - 2x^2,$$

$$2x^2 - 51x = -135;$$

$$\therefore x^2 - \frac{51}{2}x + \frac{51^2}{4} = -\frac{135}{2} + \frac{2601}{16} = \frac{1521}{16};$$

$$\therefore x - \frac{51}{4} = \pm \frac{39}{11},$$

$$\text{and } x = \text{or } \frac{45}{2}$$

$$\text{Again, } \frac{9-16x}{15x-16x^2} = \frac{32}{15}$$

$$135 - 240x = 576x - 512x^2;$$

$$\therefore 512x^2 - 516x = -135;$$

$$x^2 - \frac{51}{32}x = -\frac{135}{512};$$

$$\therefore x^2 - \frac{51}{32}x + \frac{2601}{4096} = \frac{2601}{4096} - \frac{135}{512} = \frac{2601-1080}{4096} = \frac{1521}{4096};$$

$$\therefore x - \frac{51}{64} = \pm \frac{39}{64}$$

$$\text{and } x = \frac{3}{16} \text{ or } \frac{45}{32}.$$

$$\therefore x = 3 \text{ or } \frac{45}{2} \text{ or } \frac{3}{16} \text{ or } \frac{45}{32}$$

$$y = 2 \text{ or } -\frac{135}{4} \text{ or } \frac{1}{5} \text{ or } -\frac{135}{32}$$

$$24 \quad \left. \begin{aligned} \text{Given } \sqrt{6\sqrt{x} - 4\sqrt{y}} - \frac{1}{2}\sqrt{x} &= 1 - \frac{1}{2}\sqrt{y} \\ \text{and } x - y &= 12 \end{aligned} \right\}$$

to find the values of x and y .

From the first equation,

$$\sqrt{x} - \sqrt{y} = 2\sqrt{4\sqrt{x} - 4\sqrt{y}} = 2;$$

$$\therefore 6\sqrt{x} - 4\sqrt{y} - 12\sqrt{4\sqrt{x} - 4\sqrt{y}} - 2 = 12 - 2 = 10$$

$$\text{and } \sqrt{4\sqrt{x} - 4\sqrt{y}} + 1 = \pm 2.$$

Solutions of Affected Quadratic Equations.

61

$$\therefore \sqrt{6(\sqrt{x} + \sqrt{y})} = 6 \text{ or } -18,$$

$$6(\sqrt{x} + \sqrt{y}) = 36 \text{ or } 324,$$

$$\sqrt{x} + \sqrt{y} = 6 \text{ or } 54,$$

$$\text{but } \sqrt{x} - \sqrt{y} = 2 \text{ or } \frac{2}{9};$$

$$\therefore \sqrt{x} = 4 \text{ or } \frac{244}{9}$$

$$\text{and } \sqrt{y} = 2 \text{ or } \frac{242}{9};$$

$$\therefore x = 16, \text{ or } \frac{59536}{81}$$

$$\text{and } y = 4 \text{ or } \frac{58564}{81}.$$

25. Given $y^4 - 432 = 12xy^3$
 and $y^3 = 12 + 2xy$ } to find the values
 of x and y .

$$\text{From the first equation, } x = \frac{y^4 - 432}{12y^3},$$

$$\text{and from the second, } x = \frac{y^3 - 12}{2y};$$

$$\therefore \frac{y^4 - 432}{12y^3} = \frac{y^3 - 12}{2y},$$

$$\text{and } y^4 - 432 = 6y^3 - 72y,$$

$$\text{or } y^4 + 72y = 6y^3 + 432,$$

$$\text{or } y(y^3 + 72) = 6(y^3 + 72);$$

$$\therefore y = 6,$$

$$\text{and } x = \frac{37-12}{12} = 2.$$

26. Given $\frac{4}{y^3} + \frac{4+y}{y} = \frac{8+4y}{x} + \frac{12y^2}{x^2}$ } to find the
 and $4y^3 - xy = x$

values of x and y .

From the second equation $x = \frac{4y^2}{y+1}$.

Substituting this value in the first,

$$\begin{aligned} \frac{4}{y^2} + \frac{4+y}{y} &= \frac{4 \cdot (y+2)(y+1)}{4y^2} + \frac{12y^2(y+1)^2}{16y^4} \\ &= \frac{y^2 + 3y + 2}{y^2} + \frac{3(y^2 + 2y + 1)}{4y^2}; \end{aligned}$$

$$\text{or } \frac{4}{y^2} + \frac{4}{y} + 1 = 1 + \frac{3}{y} + \frac{2}{y^2} + \frac{3}{4} + \frac{3}{2y} + \frac{3}{4y^2};$$

$$\therefore \frac{5}{4y^2} - \frac{1}{2y} = \frac{3}{4}$$

$$\text{and } 3y^2 + 2y = 5;$$

$$\therefore y^2 + \frac{2y}{3} + \frac{1}{9} = \frac{5}{3} + \frac{1}{9} = \frac{16}{9},$$

$$y + \frac{1}{3} = \pm \frac{4}{3};$$

$$\therefore y = 1 \text{ or } -\frac{5}{3},$$

$$\text{and } x = \frac{4y^2}{y+1} = 2 \text{ or } -\frac{50}{3}.$$

$$\left. \begin{aligned} 27. \quad &\text{Given } \sqrt{(1+x)^2 + y^2} + \sqrt{(1-x)^2 + y^2} = 4 \\ &\text{and } (4-x^2)^2 = 18 - 4y^2, \end{aligned} \right\}$$

to find the values of x and y .

Inverting both sides of the first equation,

$$\frac{1}{\sqrt{(1+x)^2 + y^2} + \sqrt{(1-x)^2 + y^2}} = \frac{1}{4}$$

Now multiplying numerator and denominator by

$$\sqrt{(1+x)^2 + y^2} - \sqrt{(1-x)^2 + y^2}, \text{ we have}$$

$$\sqrt{(1+x)^2 + y^2} - \sqrt{(1-x)^2 + y^2} = x,$$

$$\text{but } \sqrt{(1+x)^2 + y^2} + \sqrt{(1-x)^2 + y^2} = 4,$$

$$\begin{aligned}\therefore 2\sqrt{(1+x)^2 + y^2} &= 4 + x, \\ \text{and } 4 + 8x + 4x^2 + 4y^2 &= 16 + 8x + x^2, \\ \therefore 4y^2 &= 12 - 3x^2 = 3(4 - x^2)\end{aligned}$$

substituting in the second equation,

$$\frac{16y^4}{9} = 18 - 4y^2.$$

$$\text{Then } y^4 + \frac{9y^2}{4} = \frac{81}{8},$$

$$\therefore y^4 + \frac{9y^2}{4} + \frac{81}{64} = \frac{81 + 648}{64} = \frac{729}{64},$$

$$y^2 + \frac{9}{8} = \pm \frac{27}{8},$$

$$y^2 = \frac{9}{4} \text{ or } -\frac{9}{2},$$

$$\therefore y = \pm \frac{3}{2} \text{ or } \pm 3\sqrt{-\frac{1}{2}},$$

$$\text{and } x = \pm 1 \text{ or } \pm \sqrt{10}.$$

$$\left. \begin{aligned} 28. \text{ Given } \frac{x + \sqrt{x+y}}{x - \sqrt{x+y}} - \frac{\sqrt{x-x-y}}{\sqrt{x+x+y}} &= 2\frac{9}{40} \\ \text{and } y^2 - \sqrt{xy^2} &= \frac{4x}{9} \end{aligned} \right\}$$

to find the values of x and y .

From the first equation,

$$\frac{x+y+\sqrt{x}}{x+y-\sqrt{x}} + \frac{x+y-\sqrt{x}}{x+y+\sqrt{x}} = 2\frac{9}{40},$$

$$\text{or } \frac{2(x+y)^2 + 2x}{(x+y)^2 - x} = 2\frac{9}{40},$$

$$\therefore 2(x+y)^2 + 2x = 2\frac{9}{40} \cdot (x+y)^2 - 2\frac{9}{40}x,$$

$$\therefore \frac{9}{40}(x+y)^2 = 4\frac{9}{40}x = \frac{169}{40}x,$$

$$\text{and } \overline{x+y}^2 = \frac{169}{9}x,$$

$$\text{and } x+y = \pm \frac{13\sqrt{x}}{3}.$$

Now, from the second equation,

$$y^2 - y\sqrt{x} + \frac{x}{4} = \frac{x}{4} + \frac{4x}{9} = \frac{25x}{36},$$

$$y - \frac{\sqrt{x}}{2} = \pm \frac{5\sqrt{x}}{6},$$

$$y = \frac{4\sqrt{x}}{3} \text{ or } -\frac{\sqrt{x}}{3}.$$

$$\text{First, } x + \frac{4\sqrt{x}}{3} = \pm \frac{13\sqrt{x}}{3},$$

$$x = \frac{9\sqrt{x}}{3} \text{ or } -\frac{17\sqrt{x}}{3} = 3\sqrt{x} \text{ or } -\frac{17\sqrt{x}}{3},$$

$$\therefore \sqrt{x} = 3 \text{ or } -\frac{17}{3},$$

$$\text{and } x = 9 \text{ or } \frac{289}{9}.$$

$$\text{Again, } x - \frac{\sqrt{x}}{3} = \pm \frac{13\sqrt{x}}{3},$$

$$x = \frac{14\sqrt{x}}{3} \text{ or } -4\sqrt{x},$$

$$\sqrt{x} = \frac{14}{3} \text{ or } -4,$$

$$x = \frac{196}{9} \text{ or } 16,$$

$$\therefore x = 9 \text{ or } \frac{196}{9} \text{ or } \frac{289}{9} \text{ or } 16,$$

$$\text{and } y = 4 \text{ or } -\frac{14}{9} \text{ or } -\frac{68}{9} \text{ or } \frac{4}{3}.$$

$$29. \quad \text{Given } \left. \begin{aligned} \frac{x + \sqrt{x^2 - y^2}}{x - \sqrt{x^2 - y^2}} &= 4 \frac{1}{4} - \frac{x - \sqrt{x^2 - y^2}}{x + \sqrt{x^2 - y^2}} \\ \text{and } x \cdot x + y &= 52 - \sqrt{x^2 + xy + 4} \end{aligned} \right\}$$

to find the values of x and y .

From the first equation, we have

$$\frac{x + \sqrt{x^2 - y^2}}{x - \sqrt{x^2 - y^2}} + \frac{x - \sqrt{x^2 - y^2}}{x + \sqrt{x^2 - y^2}} = \frac{17}{4}.$$

Reducing to a common denominator,

$$\frac{2x^2 + 2x^2 - 2y^2}{y^2} = \frac{17}{4},$$

$$\therefore 4x^2 = \frac{17}{4}y^2 + 2y^2 = \frac{25y^2}{4},$$

$$\therefore 4x = 5y.$$

Again, from the second equation,

$$x^2 + xy + 4 + \sqrt{x^2 + xy + 4} + \frac{1}{4} = 52 + \frac{17}{4} = \frac{225}{4},$$

$$\sqrt{x^2 + xy + 4} + \frac{1}{2} = \pm \frac{15}{2},$$

$$\sqrt{x^2 + xy + 4} = 7 \text{ or } -8,$$

$$\therefore x^2 + xy + 4 = 49 \text{ or } 64,$$

$$x^2 + xy = 45 \text{ or } 60,$$

$$\text{but } y = \frac{4x}{5},$$

$$x^2 + \frac{4x^2}{5} = 45 \text{ or } 60,$$

$$\text{or } \frac{9x^2}{5} = 45 \text{ or } 60,$$

$$\therefore 9x^2 = 225 \text{ or } 300,$$

$$x^2 = 25 \text{ or } \frac{100}{3},$$

$$\text{and } x = \pm 5 \text{ or } \pm \frac{10}{\sqrt{3}},$$

$$\therefore y = \frac{4x}{5} = \pm 4 \text{ or } \pm \frac{8}{\sqrt{3}}.$$

$$30. \quad \text{Given } 5y + \frac{\sqrt{x^2 - 15y - 14}}{5} = \frac{x^2}{3} - 36 \quad \left\{ \begin{array}{l} \text{to find the} \\ \text{values of} \\ x \text{ and } y. \end{array} \right.$$

$$\text{and } \frac{x^2}{8y} + \frac{2x}{3} = \sqrt{\frac{x^2}{3y} + \frac{x^2}{4}} + \frac{y}{2}$$

From the first equation,

$$x^2 - 15y - 108 = \frac{3}{5} \sqrt{x^2 - 15y - 14},$$

$$\therefore x^2 - 15y - 14 - \frac{3}{5} \sqrt{x^2 - 15y - 14} + \frac{9}{100} = \frac{9}{100} + 94$$

$$= \frac{9409}{100}$$

Extracting the square root,

$$\sqrt{x^2 - 15y - 14} - \frac{3}{10} = \pm \frac{97}{10},$$

$$\therefore \sqrt{x^2 - 15y - 14} = 10 \text{ or } -\frac{47}{5}$$

$$x^2 - 15y - 14 = 100 \text{ or } \frac{2209}{25},$$

$$x^2 - 15y = 114 \text{ or } \frac{2559}{25}.$$

Again, from the second equation,

$$\frac{x^2}{8y} + \frac{2x}{3} + \frac{y}{2} = \sqrt{\frac{x^2}{3y} + \frac{x^2}{4}}.$$

Squaring both sides,

$$\frac{x^4}{64y^2} + \frac{4x^2}{9} + \frac{y^2}{4} + \frac{x^2}{6y} + \frac{x^2}{8} + \frac{2xy}{3} = \frac{x^2}{3y} + \frac{x^2}{4}$$

$$\text{Hence } \frac{x^4}{64y^2} + \frac{4x^2}{9} + \frac{y^2}{4} - \frac{x^2}{6y} - \frac{x^2}{8} + \frac{2xy}{3} = 0,$$

and extracting the square root,

$$\frac{x^2}{8y} - \frac{2x}{3} - \frac{y}{2} = 0,$$

$$\therefore x^2 - \frac{16xy}{3} = 4y^2,$$

$$x^2 - \frac{16xy}{3} + \frac{64y^2}{9} = \frac{(36 - 64)y^2}{9} = \frac{100y^2}{9}$$

$$x - \frac{8y}{3} = \pm \frac{10y}{3}$$

$$\text{and } x = 8y \text{ or } -\frac{2y}{3}$$

$$y = \frac{x}{6} \text{ or } -\frac{3x}{2}$$

$$\text{First, } x^2 - 15 \cdot \frac{x}{6} = 114 \text{ or } \frac{2339}{25}$$

$$\text{or } x^2 - \frac{5x}{2} + \frac{25}{16} = \frac{25}{16} + 114 = \frac{1849}{16} \text{ or } \frac{41569}{400}$$

$$\therefore x - \frac{5}{4} = \pm \frac{43}{4} \text{ or } \pm \frac{\sqrt{41569}}{20}$$

$$\text{and } x = 12 \text{ or } -\frac{19}{2} \text{ or } \frac{5}{4} \pm \frac{\sqrt{41569}}{20}$$

$$= 12 \text{ or } -\frac{19}{2} \text{ or } \frac{25 \pm \sqrt{41569}}{20}.$$

$$\text{Again, } x^2 - 15 \left(-\frac{3x}{2}\right) = 114,$$

$$\text{or } x^2 + \frac{45x}{2} + \left(\frac{45}{4}\right)^2 = 114 + \frac{2025}{16} = \frac{3849}{16},$$

$$x + \frac{45}{4} = \pm \frac{\sqrt{3849}}{4},$$

$$\text{and } x = \frac{-45 \pm \sqrt{3849}}{4}.$$

$$\therefore x = 12 \text{ or } -\frac{19}{2} \text{ or } \frac{25 \pm \sqrt{41569}}{20} \text{ or } \frac{-45 \pm \sqrt{3849}}{4},$$

$$\text{and } y = 2 \text{ or } -\frac{19}{2} \text{ or } \frac{25 \pm \sqrt{41569}}{120} \text{ or } \frac{-135 \pm \sqrt{3849}}{8}.$$

$$31. \quad \left. \begin{aligned} \text{Given } \sqrt{\frac{x+y^2}{4x}} + \frac{y}{\sqrt{y^2+x}} &= \frac{y^2}{4} \sqrt{\frac{4x}{y^2+x}} \\ \text{and } \frac{\sqrt{x} + \sqrt{x-y-1}}{\sqrt{x} - \sqrt{x-y-1}} &= y+1 \end{aligned} \right\}$$

to find the values of x and y .

Multiplying the first equation by $\sqrt{\frac{x+y^2}{4x}}$,

$$\frac{x+y^2}{4x} + \frac{y}{\sqrt{4x}} = \frac{y^2}{4},$$

$$\text{or } \frac{y^2}{x} + \frac{2y}{\sqrt{x}} + 1 = y^2,$$

$$\text{and } \frac{y}{\sqrt{x}} + 1 = \pm y,$$

$$\pm \sqrt{x} y - \sqrt{x} = y,$$

$$\sqrt{x} = \frac{y}{\pm y - 1}.$$

Multiplying numerator and denominator of the left side of the second equation by $\sqrt{x} + \sqrt{x-y-1}$,

$$\frac{2x - y - 1 + 2\sqrt{x}\sqrt{x-y-1}}{y+1} = y+1,$$

$$\text{and } 2x - y - 1 + 2\sqrt{x}\sqrt{x-y-1} = (y+x)^2 \quad (1).$$

$$\text{Again, } \frac{\sqrt{x} - \sqrt{x-y-1}}{\sqrt{x} + \sqrt{x-y-1}} = \frac{1}{y+1},$$

and multiplying numerator and denominator as before by

$$\sqrt{x} - \sqrt{x-y-1}, \text{ we have}$$

$$2x - y - 1 - 2\sqrt{x}\sqrt{x-y-1} = 1 \quad (2).$$

adding together equations (1) and (2)

$$4x - 2y - 2 = 1 + y + 1)^2$$

$$\text{and } 4x = y^2 + 4y + 4,$$

$$\therefore 2\sqrt{x} = y + 2,$$

$$\text{but } 2\sqrt{x} = \frac{2y}{\pm y - 1}$$

$$\therefore y + 2 = \frac{2y}{\pm y - 1}$$

$$\text{and first } y + 2 = \frac{2y}{y - 1}$$

$$\therefore y^2 + y - 2 = 2y,$$

$$y^2 - y + \frac{1}{4} = \frac{9}{4},$$

$$y - \frac{1}{2} = \pm \frac{3}{2},$$

$$y = 2 \text{ or } -1.$$

$$\text{Again, } y + 2 = \frac{2y}{-y - 1},$$

$$\text{or } -y^2 - 3y - 2 = 2y,$$

$$y^2 + 5y + \frac{25}{4} = \frac{25}{4} - 2 = \frac{17}{4},$$

$$\therefore y + \frac{5}{2} = \pm \frac{\sqrt{17}}{2},$$

$$y = \frac{-5 \pm \sqrt{17}}{2},$$

$$\therefore y = 2, \text{ or } -1, \text{ or } \frac{-5 \pm \sqrt{17}}{2},$$

$$\text{and } x = 4 \text{ or } \frac{1}{4} \text{ or } \frac{13 \pm \sqrt{17}}{8}.$$

$$32. \quad \text{Given } \left. \begin{aligned} \frac{x + y + \sqrt{x^2 - y^2}}{x + y - \sqrt{x^2 - y^2}} &= \frac{9}{8y} (x + y) \\ \text{and } (x^2 + y)^2 + x - y &= 2x(x^2 + y) + 506 \end{aligned} \right\} \begin{array}{l} \text{to find the va-} \\ \text{lues of } x \\ \text{and } y. \end{array}$$

Dividing numerator and denominator of the left hand-side of the first equation by $\sqrt{x + y}$,

$$\frac{\sqrt{x + y} + \sqrt{x - y}}{\sqrt{x + y} - \sqrt{x - y}} = \frac{9}{8y} (x + y).$$

Again, multiplying numerator and denominator by

$$\frac{\sqrt{x+y} + \sqrt{x-y}}{2y} = \frac{9}{8y}(x+y)$$

$$\text{or } (\sqrt{x+y} + \sqrt{x-y})^2 = \frac{9}{4}(x+y)$$

$$\text{and } \therefore \sqrt{x+y} + \sqrt{x-y} = \frac{3}{2}\sqrt{x+y},$$

$$\therefore \frac{\sqrt{x+y}}{2} = \sqrt{x-y}.$$

Squaring both sides, and clearing of fractions,

$$x+y = 4x-4y$$

$$3x = 5y$$

$$y = \frac{3x}{5}.$$

Again, from the second equation,

$$(x^2 + y)^2 - 2x(x^2 + y) + x^2 = x^2 + y - x + 506,$$

$$(x^2 + y - x)^2 - (x^2 + y - x) + \frac{1}{4} = 506 + \frac{1}{4} = \frac{2025}{4}$$

$$x^2 + y - x = \frac{\pm 45 + 1}{2} = 23 \text{ or } -22,$$

$$\text{or } x^2 + \frac{3x}{5} - x = 23 \text{ or } -22,$$

$$x^2 - \frac{2x}{5} + \frac{1}{25} = 23 + \frac{1}{25} \text{ or } -22 + \frac{1}{25}$$

$$= \frac{576}{25} \text{ or } -\frac{549}{25},$$

$$x - \frac{1}{5} = \pm \frac{24}{5} \text{ or } \pm \frac{\sqrt{-549}}{5},$$

$$x = 5 \text{ or } -\frac{23}{5} \text{ or } \frac{1 \pm \sqrt{-549}}{5},$$

$$y = 3 \text{ or } -\frac{69}{25} \text{ or } \frac{3 \pm 3\sqrt{-549}}{25}.$$

$$33. \quad \left. \begin{aligned} \text{Given } \frac{y}{x} \cdot \sqrt{\frac{x}{y}} + \frac{1}{2} \sqrt{\frac{x}{y}} \cdot \sqrt{\frac{y^3}{x^3}} &= 5 \\ \text{and } \frac{2x^2}{y} - \frac{x}{3\sqrt{y}} &= \frac{1}{3} \end{aligned} \right\}$$

to find the values of x and y .

From the first equation

$$\sqrt{\frac{y}{x}} + \frac{1}{2} \cdot \frac{x^{\frac{1}{2}}y^{\frac{3}{2}}}{y^{\frac{1}{2}}x^{\frac{3}{2}}} = 5,$$

$$\text{or } \frac{y^{\frac{1}{2}}}{x^{\frac{1}{2}}} + \frac{1}{2} \cdot \frac{y^{\frac{1}{2}}}{x^{\frac{1}{2}}} + \frac{1}{16} = \frac{1}{16} + 5 = \frac{81}{16},$$

$$\frac{y^{\frac{1}{2}}}{x^{\frac{1}{2}}} + \frac{1}{4} = \pm \frac{9}{4},$$

$$\frac{y^{\frac{1}{2}}}{x^{\frac{1}{2}}} = 2 \text{ or } -\frac{5}{2},$$

$$\frac{y}{x} = 16 \text{ or } \frac{625}{16}.$$

From the second equation,

$$\frac{x^2}{y} - \frac{1}{6} \cdot \frac{x}{\sqrt{y}} + \frac{1}{144} = \frac{1}{144} + \frac{1}{6} = \frac{25}{144}$$

$$\therefore \frac{x}{\sqrt{y}} - \frac{1}{12} = \pm \frac{5}{12},$$

$$\frac{x}{\sqrt{y}} = \frac{1}{2} \text{ or } -\frac{1}{2},$$

$$\text{and } \frac{x^2}{y} = \frac{1}{4} \text{ or } \frac{1}{9},$$

$$\therefore \frac{x^2}{y} \times \frac{y}{x} = \frac{1}{4} \times 16 \text{ or } \frac{1}{4} \times \frac{625}{16} \text{ or } \frac{1}{9} \times 16 \text{ or } \frac{1}{9} \times \frac{625}{16},$$

$$\therefore x = 4, \text{ or } \frac{625}{64}, \text{ or } \frac{16}{9}, \text{ or } \frac{625}{144},$$

$$\text{and } y = 64 \text{ or } \frac{625}{32} \text{ or } \frac{256}{9} \text{ or } \frac{625}{45}.$$

$$34. \quad \left. \begin{array}{l} \text{Given } \sqrt{\frac{x}{y}} + \sqrt{\frac{y}{x}} = \frac{61}{\sqrt{xy}} + 1 \\ \text{and } \sqrt[4]{x^3y} + \sqrt[4]{y^3x} = 78 \end{array} \right\} \begin{array}{l} \text{to find the va-} \\ \text{lues of } x \\ \text{and } y. \end{array}$$

From the first equation,

$$\frac{x+y}{\sqrt{xy}} = \frac{61}{\sqrt{xy}} + 1,$$

$$\therefore x+y = 61 + \sqrt{xy}.$$

From the second,

$$(\sqrt{xy})^{\frac{1}{2}} (\sqrt{x} + \sqrt{y}) = 78,$$

$$\text{or } \sqrt{x} + \sqrt{y} = \frac{78}{\sqrt[4]{xy}},$$

$$\therefore x+y+2\sqrt{xy} = \frac{6084}{\sqrt{xy}},$$

$$\text{and } \therefore x+y = \frac{6084}{\sqrt{xy}} - 2\sqrt{xy} = 61 + \sqrt{xy},$$

$$\therefore 6084 = 61\sqrt{xy} + 3xy,$$

$$\text{and } xy + \frac{61}{3}\sqrt{xy} + \frac{61}{6} = \frac{6084}{3} + \frac{61}{6} \Rightarrow \frac{76729}{36},$$

$$\therefore \sqrt{xy} + \frac{61}{6} = \pm \frac{277}{6}$$

$$\text{and } \sqrt{xy} = \frac{216}{6} \text{ or } -\frac{338}{6} = 36 \text{ or } -\frac{169}{3},$$

$$xy = 1296 \text{ or } \frac{28561}{9},$$

$$\text{but } x+y = 61 + \sqrt{xy} = 97,$$

$$\therefore x^2 + 2xy + y^2 = 9409 \text{ or } \frac{196}{9},$$

$$\text{but } 4xy = 5184 \text{ or } \frac{114244}{9},$$

$$x^2 - 2xy + y^2 = 4225 \text{ or } -\frac{114048}{9},$$

$$x - y = \pm 65 \text{ or } \pm 24\sqrt{-22}$$

$$\text{but } x + y = 97$$

$$x = 81, \text{ or } 16, \text{ or } \frac{97}{2} \pm 12\sqrt{-22},$$

$$y = 16 \text{ or } 81, \text{ or } \frac{97}{2} \mp 12\sqrt{-22}.$$

$$35. \text{ Given } \left. \begin{aligned} \frac{y}{2x} + \frac{2}{3} \cdot \frac{y - \sqrt{x-1}}{y^2 - 2\sqrt{x^2-1}} &= \frac{\sqrt{x+1}}{x} \\ \text{and } \frac{1}{4} \cdot y^4 &= y^2x - 1, \end{aligned} \right\}$$

to find the values of x and y .

From the second equation,

$$y^4 = 4y^2x - 4,$$

$$\therefore y^4 - 4y^2x + 4x^2 = 4(x^2 - 1),$$

$$\therefore y^2 - 2x = \pm 2\sqrt{x^2 - 1};$$

$$\begin{aligned} \therefore y^2 &= 2x \pm 2\sqrt{x^2 - 1} \\ &= (\sqrt{x+1} \pm \sqrt{x-1})^2 \end{aligned}$$

$$\text{and } y = \sqrt{x+1} \pm \sqrt{x-1}.$$

Substituting for $2\sqrt{x^2 - 1}$, $y^2 - 2x$ in the second equation,

$$\frac{y}{2x} + \frac{2}{3} \cdot \frac{y - \sqrt{x-1}}{2x} = \frac{\sqrt{x+1}}{x},$$

$$\text{or } \frac{y}{2} + \frac{1}{3}(y - \sqrt{x-1}) = \sqrt{x+1}.$$

Substituting for y , $\sqrt{x+1} + \sqrt{x-1}$

$$\frac{y}{2} + \frac{1}{3}\sqrt{x+1} = \sqrt{x+1},$$

$$\frac{y}{2} = \frac{2\sqrt{x+1}}{3} = \frac{\sqrt{x+1} + \sqrt{x-1}}{2};$$

$$\therefore \frac{\sqrt{x+1}}{3} = \sqrt{x-1}, \text{ and } x+1 = 9x-9;$$

$$\therefore 8x = 10 \text{ and } x = \frac{5}{4}.$$

Again, substituting for y , $\sqrt{x+1} - \sqrt{x-1}$

$$\frac{y}{2} + \frac{1}{3}(\sqrt{x+1} - 2\sqrt{x-1}) = \sqrt{x+1}$$

$$\frac{y}{2} = \frac{2}{3}(\sqrt{x-1} + \sqrt{x+1}) = \frac{\sqrt{x+1} - \sqrt{x-1}}{2}$$

$$4\sqrt{x+1} + 4\sqrt{x-1} = 3\sqrt{x+1} - 3\sqrt{x-1}$$

$$\therefore \sqrt{x+1} = -7\sqrt{x-1}$$

$$x+1 = 49x-49,$$

$$48x = 50,$$

$$\text{and } x = \frac{50}{48} = \frac{25}{24};$$

$$\therefore x = \frac{5}{4} \text{ or } \frac{25}{24}.$$

$$\text{Now } y^4 - 4y^2x^2 = -4,$$

$$\text{or } y^4 - 5y^2 = -4,$$

$$y^4 - 5y^2 + \frac{25}{4} = \frac{9}{4},$$

$$y^2 - \frac{5}{2} = \pm \frac{3}{2},$$

$$y^2 = 4 \text{ or } 1,$$

$$y = \pm 2 \text{ or } \pm 1.$$

$$\text{Again, } y^4 - \frac{25y^2}{6} = -4,$$

$$y^4 - \frac{25y^2}{6} + \frac{625}{144} = \frac{625-576}{144} = \frac{49}{144};$$

$$\therefore y^2 - \frac{25}{12} = \pm \frac{7}{12},$$

$$y^2 = \frac{8}{3} \text{ or } \frac{3}{2};$$

$$\therefore y = \pm \frac{2}{3} \sqrt{6} \text{ or } \pm \frac{1}{2} \sqrt{6}.$$

36. Given $x^2 - y^2 = 3$,
 and $(x^4 + y^4)^2 + x^2y^2(x^2 - y^2)^2 + x^2 - y^2 = 328$ } to find
 the values of x and y .

Substituting 3 for $x^2 - y^2$ in the second equation,

$$(x^4 + y^4)^2 + 9x^2y^2 + 3 = 328,$$

$$\text{and } x^4 - 2x^2y^2 + y^4 = 9;$$

$$\therefore x^4 + y^4 = 9 + 2x^2y^2;$$

$$\therefore \text{also } (9 + 2x^2y^2)^2 + 9x^2y^2 = 325,$$

$$\text{or } 4x^4y^4 + 45x^2y^2 = 244;$$

$$\therefore x^4y^4 + \frac{45}{4}x^2y^2 + \left(\frac{45}{8}\right)^2 = \left(\frac{45}{8}\right)^2 + \frac{244}{4} = \frac{5929}{64};$$

$$\therefore x^2y^2 + \frac{45}{8} = \pm \frac{77}{8},$$

$$\text{and } x^2y^2 = 4 \text{ or } -\frac{61}{4}.$$

$$\text{Now } x^4 - 2x^2y^2 + y^4 = 9,$$

$$4x^2y^2 = 16 \text{ or } -61.$$

$$x^4 + 2x^2y^2 + y^4 = 25 \text{ or } -52$$

$$\text{and } x^2 + y^2 = \pm 5 \text{ or } \pm 2\sqrt{-13},$$

$$\text{but } x^2 - y^2 = 3;$$

$$\therefore x^2 = 4 \text{ or } -1 \text{ or } \frac{3 \pm 2\sqrt{-13}}{2},$$

$$\text{and } x = \pm 2 \text{ or } \pm \sqrt{-1} \text{ or } \pm \sqrt{\frac{3 \pm 2\sqrt{-13}}{2}}.$$

$$\text{Similarly } y = \pm 1 \text{ or } \pm 2\sqrt{-1} \text{ or } \pm \sqrt{\frac{3 \mp 2\sqrt{-13}}{2}}.$$

$$37. \text{ Given } \frac{2y^2 - 8\sqrt{x}}{\sqrt{x}} + \frac{\sqrt{4y^2 - 16\sqrt{x}}}{2} = \frac{3\sqrt{x}}{2} \}$$

$$\text{and } \sqrt{x} + \sqrt{8(y - \sqrt{x}) - 4} = y + 1$$

to find the values of x and y .

From the first equation,

$$y^2 - 4\sqrt{x} + \sqrt{x}\sqrt{y^2 - 4\sqrt{x}} = \frac{3x}{4}$$

$$\therefore y^2 - 4\sqrt{x} + \sqrt{x}\sqrt{y^2 - 4\sqrt{x}} + \frac{x}{4} = \frac{3x}{4} + \frac{x}{4} = x;$$

$$\therefore \sqrt{y^2 - 4\sqrt{x}} + \frac{\sqrt{x}}{2} = \pm \sqrt{x},$$

$$\sqrt{y^2 - 4\sqrt{x}} = \frac{\sqrt{x}}{2} \text{ or } -\frac{3\sqrt{x}}{2}$$

$$y^2 - 4\sqrt{x} = \frac{x}{4} \text{ or } \frac{9x}{4},$$

$$\text{and } y^2 = \frac{x}{4} + 4\sqrt{x}, \text{ or } \frac{9x}{4} + 4\sqrt{x}.$$

Again, from the second equation,

$$y - \sqrt{x} + 1 = 2\sqrt{2(y - \sqrt{x}) - 1};$$

$$\therefore 2(y - \sqrt{x}) + 2 = 4\sqrt{2(y - \sqrt{x}) - 1},$$

$$2(y - \sqrt{x}) - 1 - 4\sqrt{2(y - \sqrt{x}) - 1} = -3;$$

$$\therefore (2(y - \sqrt{x}) - 1) - 4\sqrt{2(y - \sqrt{x}) - 1} + 4 = 1,$$

$$\sqrt{2(y - \sqrt{x}) - 1} - 2 = \pm 1,$$

$$\sqrt{2(y - \sqrt{x}) - 1} = 3 \text{ or } 1,$$

$$2(y - \sqrt{x}) - 1 = 9 \text{ or } 1.$$

$$\therefore y - \sqrt{x} = 5 \text{ or } 1,$$

$$y = 5 + \sqrt{x}, \text{ or } 1 + \sqrt{x};$$

$$\therefore y^2 = x + 10\sqrt{x} + 25, \text{ or } x + 2\sqrt{x} + 1,$$

$$\text{but } y^2 = \frac{x}{4} + 4\sqrt{x}, \text{ or } \frac{9x}{4} + 4\sqrt{x}.$$

First then,

$$x + 2\sqrt{x} + 1 = \frac{x}{4} + 4\sqrt{x} \text{ or } \frac{9x}{4} + 4\sqrt{x},$$

$$\text{or } x - 2\sqrt{x} + 1 = \frac{x}{4} \text{ or } \frac{9x}{4},$$

$$\sqrt{x} - 1 = \pm \frac{\sqrt{x}}{2} \text{ or } -\frac{3\sqrt{x}}{2};$$

$$\frac{\sqrt{x}}{2} = 1 \text{ or } \frac{3\sqrt{x}}{2} = 1 \text{ or } 5\sqrt{x} = 2;$$

$$\therefore \sqrt{x} = 2 \text{ or } \frac{2}{3} \text{ or } \sqrt{x} = \frac{2}{5};$$

$$\therefore x = 4 \text{ or } \frac{4}{9}, \text{ and } x = \frac{4}{25}.$$

Secondly,

$$x + 10\sqrt{x} + 25 = \frac{x}{4} + 4\sqrt{x};$$

$$\therefore \frac{3x}{4} + 6\sqrt{x} = -25,$$

$$x + 8\sqrt{x} + 16 = 16 - \frac{100}{3} = -\frac{52}{3},$$

$$\sqrt{x} + 4 = \pm \sqrt{-\frac{52}{3}} = \pm 2\sqrt{-\frac{13}{3}},$$

$$\sqrt{x} = -4 \pm 2\sqrt{-\frac{13}{3}},$$

$$\text{and } x = 16 \pm 16\sqrt{-\frac{13}{3}} - \frac{52}{3} = -\frac{4}{3} \mp 16\sqrt{-\frac{13}{3}}.$$

Thirdly,

$$x + 10\sqrt{x} + 25 = \frac{9x}{4} + 4\sqrt{x};$$

$$\therefore \frac{5x}{4} - 6\sqrt{x} = 25,$$

$$x - \frac{24\sqrt{x}}{5} = 20;$$

$$\therefore x - \frac{24\sqrt{x}}{5} + \frac{144}{25} = 20 + \frac{144}{25} = \frac{644}{25},$$

$$\sqrt{x} - \frac{12}{5} = \pm \frac{\sqrt{644}}{25},$$

$$\sqrt{x} = \frac{12 \pm \sqrt{644}}{25},$$

$$\text{and } x = \frac{786 \pm 24\sqrt{644}}{25};$$

$$\therefore x = 4, \text{ or } \frac{4}{9}; \text{ or } \frac{4}{25}, \text{ or } -\frac{4}{3} \mp 16\sqrt{-\frac{13}{3}},$$

$$\text{or } \frac{788 \pm 24\sqrt{644}}{25},$$

$$\text{and } y = 3, \text{ or } \frac{5}{3}; \text{ or } \frac{7}{5}, \text{ or } -1 \text{ or } 1 \pm 2\sqrt{-\frac{13}{3}}$$


$$\text{or } \frac{37 \pm \sqrt{644}}{5}.$$

THE END.

Shortly will be published, by the same Author,
**REMARKS UPON THE PRESENT SYSTEM OF
EDUCATION AT PRIVATE SCHOOLS;
WITH PARTICULAR REFERENCE TO THEIR CONDUCTORS.**

Dedicated (by permission)
TO THE RIGHT HONOURABLE JOHN, EARL OF ELDON.

LONDON :
Printed by WILLIAM CLOWES,
Stamford-street.





•

•

•



